



# Need more money for college expenses?

The CLC Private Loan<sup>SM</sup> can get you up to  
**\$40,000** a year for college-related expenses.

---

Here's why the CLC Private Loan<sup>SM</sup> is a smart choice:

- ✓ Approved borrowers are sent a check within four business days
- ✓ Get \$1000 - \$40,000 each year
- ✓ Interest rates as low as prime + 0% (8.66% APR)
- ✓ Quick and easy approval process
- ✓ No payments until after graduating or leaving school

**CLICK HERE**

or call **800.311.9615.**

*We are available 24 hours  
a day, 7 days a week.*

# Operations with Rational Expressions

Marc owns  $\frac{5}{8}$  interest in a print shop and his uncle owns  $\frac{1}{4}$  interest in the shop. In a given year, they shared earnings of \$140,000. How much did the shop earn that year?



## 6-1 ■ Multiplication and division of rational expressions

### Multiplication of rational expressions

Recall that to multiply two real number fractions we multiply the numerators and multiply the denominators.

#### Multiplication property of fractions

If  $a$ ,  $b$ ,  $c$ , and  $d$  are real numbers, then

$$\frac{a}{b} \cdot \frac{c}{d} = \frac{a \cdot c}{b \cdot d} \quad (b, d \neq 0)$$

**Note** Any possible reduction is performed *before* the multiplication takes place.

### ■ Example 6-1 A

Multiply the fractions  $\frac{3}{7}$  and  $\frac{14}{27}$  and simplify the product.

$$\begin{aligned} \frac{3}{7} \cdot \frac{14}{27} &= \frac{3 \cdot 14}{7 \cdot 27} \\ &= \frac{3 \cdot 2 \cdot 7}{7 \cdot 3 \cdot 3 \cdot 3} \\ &= \frac{2 \cdot (3 \cdot 7)}{3 \cdot 3 \cdot (3 \cdot 7)} \end{aligned}$$

Multiply the numerators  
Multiply the denominators

Factor the numerator and the denominator

Group the common factors ( $3 \cdot 7$ )



$$= \frac{2}{3 \cdot 3}$$

$$= \frac{2}{9}$$

Divide numerator and denominator by  $(3 \cdot 3)$ 

Multiply the remaining factors

This same procedure is followed when we multiply two rational expressions.

### Multiplication property of rational expressions

Given rational expressions  $\frac{P}{Q}$  and  $\frac{R}{S}$ , then

$$\frac{P}{Q} \cdot \frac{R}{S} = \frac{P \cdot R}{Q \cdot S} \quad (Q, S \neq 0)$$

Since we want the resulting product to be stated in lowest terms, we apply the *fundamental principle of rational expressions* and divide both the numerator and the denominator by their common factors. That is, we *reduce* by dividing out the common factors.

### Multiplication of rational expressions

1. State the numerators and denominators as indicated products. (Do not multiply.)
2. Factor the numerator and the denominator.
3. Divide the numerators and the denominators by the factors that are common.
4. Multiply the remaining factors in the numerator and place this product over the product of the remaining factors in the denominator.

### Example 6-1 B

Perform the indicated multiplication and simplify your answer. Assume that no denominator equals zero.

$$1. \frac{3x}{4} \cdot \frac{5}{2y}$$

$$= \frac{3x \cdot 5}{4 \cdot 2y}$$

$$= \frac{15x}{8y}$$

Multiply numerators and denominators

Will not reduce

$$2. \frac{x+1}{x-3} \cdot \frac{4}{x+2}$$

$$= \frac{(x+1) \cdot 4}{(x-3)(x+2)}$$

$$= \frac{4x+4}{x^2-x-6}$$

Multiply numerators and denominators

Will not reduce

$$\begin{aligned}
 3. \quad & \frac{4}{9x} \cdot \frac{3x^2}{2} \\
 &= \frac{4 \cdot 3x^2}{9x \cdot 2} \\
 &= \frac{2 \cdot 2 \cdot 3 \cdot x \cdot x}{3 \cdot 3 \cdot 2 \cdot x} \\
 &= \frac{2 \cdot x \cdot (2 \cdot 3 \cdot x)}{3 \cdot (2 \cdot 3 \cdot x)} \\
 &= \frac{2x}{3}
 \end{aligned}$$

Multiply numerators

Multiply denominators

Factor numerator and denominator

Identify common factors

Divide numerator and denominator by  $(2 \cdot 3 \cdot x)$ 

$$\begin{aligned}
 4. \quad & \frac{x+1}{3-x} \cdot \frac{(x-3)^2}{x-2} \\
 &= \frac{(x+1) \cdot (x-3)^2}{(3-x) \cdot (x-2)} \\
 &= \frac{x-3}{3-x} \cdot \frac{(x+1)(x-3)}{x-2} \\
 &= \frac{-1 \cdot (x+1)(x-3)}{x-2} \\
 &= \frac{-1(x^2 - 2x - 3)}{x-2} \\
 &= \frac{-x^2 + 2x + 3}{x-2}
 \end{aligned}$$

Multiply numerators

Multiply denominators

Factor opposites

$$\frac{x-3}{3-x} = -1$$

Multiply as indicated

$$\begin{aligned}
 5. \quad & \frac{x^2 - 8x + 16}{x^2 + 3x - 10} \cdot \frac{x^2 - 4}{x^2 - 5x + 4} \\
 &= \frac{(x^2 - 8x + 16)(x^2 - 4)}{(x^2 + 3x - 10)(x^2 - 5x + 4)} \\
 &= \frac{(x-4)(x-4)(x-2)(x+2)}{(x-4)(x-2)(x+5)(x-1)} \\
 &= \frac{(x-4)(x+2)}{(x+5)(x-1)} \\
 &= \frac{x^2 - 2x - 8}{x^2 + 4x - 5}
 \end{aligned}$$

Multiply numerators and denominators

Factor numerator and denominator

Reduce by common factors  $(x-4)$  and  $(x-2)$ 

Multiply remaining factors

► **Quick check** Multiply  $\frac{12}{5y} \cdot \frac{15y^2}{4}$

### Division of rational expressions

Recall that to divide two fractions  $\frac{a}{b}$  and  $\frac{c}{d}$ , we multiply  $\frac{a}{b}$  by the *reciprocal* of

$\frac{c}{d}$ , which is  $\frac{d}{c}$ .



**Division property of fractions**If  $a$ ,  $b$ ,  $c$ , and  $d$  are real numbers, then

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c} = \frac{a \cdot d}{b \cdot c} \quad (b, c, d \neq 0)$$

**Example 6-1 C**

Find the indicated quotient. Reduce your answer to lowest terms.

$$\begin{aligned} \frac{18}{25} \div \frac{9}{5} &= \frac{18}{25} \cdot \frac{5}{9} && \text{Multiply by the reciprocal of } \frac{9}{5} \\ &= \frac{2 \cdot 3 \cdot 3 \cdot 5}{5 \cdot 5 \cdot 3 \cdot 3} && \text{Factor in numerator and denominator} \\ &= \frac{2 \cdot (3 \cdot 3 \cdot 5)}{5 \cdot (3 \cdot 3 \cdot 5)} && \text{Group the common factors } (3 \cdot 3 \cdot 5) \\ &= \frac{2}{5} && \text{Reduce by the common factors } (3 \cdot 3 \cdot 5) \end{aligned}$$

Division of rational expressions is done in the same way.

**Division property of rational expressions**If  $\frac{P}{Q}$  and  $\frac{R}{S}$  are rational expressions, then

$$\frac{P}{Q} \div \frac{R}{S} = \frac{P}{Q} \cdot \frac{S}{R} = \frac{P \cdot S}{Q \cdot R} \quad (Q, R, S \neq 0)$$

Notice that once the operation of division has been changed to multiplication, we proceed exactly as we did with the multiplication of rational expressions.

**Division of rational expressions**

1. Multiply the first rational expression by the reciprocal of the second.
2. Proceed as in the multiplication of rational expressions.

**Example 6-1 D**

Find the indicated quotients. Express the answer in reduced form.

$$\begin{aligned} 1. \quad \frac{3ab}{5} \div \frac{9abc}{10} &= \frac{3ab}{5} \cdot \frac{10}{9abc} && \text{Multiply by the reciprocal of } \frac{9abc}{10} \\ &= \frac{3ab \cdot 2 \cdot 5}{5 \cdot 3 \cdot 3 \cdot abc} && \text{Factor numerator and denominator} \\ &= \frac{2 \cdot (3 \cdot 5 \cdot ab)}{3 \cdot c \cdot (3 \cdot 5 \cdot ab)} && \text{Group common factors } (3 \cdot 5 \cdot ab) \\ &= \frac{2}{3c} && \text{Divide numerator and denominator by common factors } (3 \cdot 5 \cdot ab) \end{aligned}$$

$$\begin{aligned}
 2. \quad \frac{x^2 - 4}{5} \div \frac{x - 2}{15} &= \frac{x^2 - 4}{5} \cdot \frac{15}{x - 2} \\
 &= \frac{(x - 2)(x + 2) \cdot 3 \cdot 5}{5 \cdot (x - 2)} \\
 &= \frac{3 \cdot (x + 2) \cdot 5(x - 2)}{5(x - 2)} \\
 &= \frac{3(x + 2)}{1} = 3x + 6
 \end{aligned}$$

Multiply by the reciprocal of  $\frac{x - 2}{15}$

Factor numerator and denominator

Locate common factors  $5(x - 2)$

Reduce to lowest terms by dividing numerator and denominator by  $5(x - 2)$

$$\begin{aligned}
 3. \quad \frac{4x + 2}{x - 1} \div \frac{2x + 1}{4 - 4x} &= \frac{4x + 2}{x - 1} \cdot \frac{4 - 4x}{2x + 1} \\
 &= \frac{2(2x + 1)(-4)(x - 1)}{(x - 1)(2x + 1)} \\
 &= \frac{2(-4)(2x + 1)(x - 1)}{(2x + 1)(x - 1)} \\
 &= 2(-4) \\
 &= -8
 \end{aligned}$$

Multiply by the reciprocal of  $\frac{2x + 1}{4 - 4x}$

Factor  $4 - 4x = -4(x - 1)$  and  $4x + 2 = 2(2x + 1)$

Reduce by  $(x - 1)(2x + 1)$

$$\begin{aligned}
 4. \quad \frac{x^2 - 9}{2x + 1} \div \frac{3 - x}{2x^2 + 7x + 3} &= \frac{x^2 - 9}{2x + 1} \cdot \frac{2x^2 + 7x + 3}{3 - x} \\
 &= \frac{(x - 3)(x + 3) \cdot (2x + 1)(x + 3)}{(2x + 1)(3 - x)} \\
 &= \frac{x - 3}{3 - x} \cdot \frac{(2x + 1)(x + 3)(x + 3)}{2x + 1} \\
 &= -1 \cdot (x + 3)(x + 3) \\
 &= -1(x^2 + 6x + 9) \\
 &= -x^2 - 6x - 9
 \end{aligned}$$

Multiply by the reciprocal of  $\frac{3 - x}{2x^2 + 7x + 3}$

Factor numerators

Factor opposites

$\frac{x - 3}{3 - x} = -1$ ; reduce by  $(2x + 1)$

Multiply as indicated

Multiply by  $-1$

► **Quick check** Divide  $\frac{x^2 - 4}{4x - 1} \div \frac{2 - x}{4x^2 + 3x - 1}$  ■

### Mastery points

Can you

- Multiply rational expressions?
- Divide rational expressions?

## Exercise 6-1

Find the indicated product or quotient. Write your answer in simplest form. Assume all denominators are nonzero. See examples 6-1 A-D.

**Example**  $\frac{12}{5y} \cdot \frac{15y^2}{4}$

**Solution** 
$$= \frac{12 \cdot 15y^2}{5y \cdot 4}$$
 Multiply numerator and denominator

$$= \frac{3 \cdot 2 \cdot 2 \cdot 3 \cdot 5 \cdot y \cdot y}{5 \cdot y \cdot 2 \cdot 2}$$
 Factor numerator and denominator
$$= \frac{3 \cdot 3 \cdot y(5 \cdot y \cdot 2 \cdot 2)}{1(5 \cdot y \cdot 2 \cdot 2)}$$
 Group common factors ( $5 \cdot y \cdot 2 \cdot 2$ )
$$= \frac{9y}{1}$$
 Divide by common factors ( $5 \cdot y \cdot 2 \cdot 2$ )
$$= 9y$$

**Example**  $\frac{x^2 - 4}{4x - 1} \div \frac{2 - x}{4x^2 + 3x - 1}$

**Solution** 
$$= \frac{x^2 - 4}{4x - 1} \cdot \frac{4x^2 + 3x - 1}{2 - x}$$
 Multiply by the reciprocal of  $\frac{2 - x}{4x^2 + 3x - 1}$

$$= \frac{(x + 2)(x - 2) \cdot (4x - 1)(x + 1)}{(4x - 1)(2 - x)}$$
 Factor numerator and denominator
$$= \frac{x - 2}{2 - x} \cdot \frac{(x + 2)(x + 1)(4x - 1)}{4x - 1}$$
 Factor opposites
$$= -1 \cdot (x + 2)(x + 1)$$
  $\frac{x - 2}{2 - x} = -1$ ; reduce by  $4x - 1$ 

$$= -x^2 - 3x - 2$$
 Multiply as indicated

1.  $\frac{24}{35} \cdot \frac{7}{8}$

5.  $\frac{4a}{5} \cdot \frac{5}{2}$

9.  $\frac{5}{6} \cdot \frac{3x}{10y}$

13.  $\frac{24a}{35x} \div 6a$

17.  $\frac{21ab}{16c} \cdot \frac{8c^2}{3ab^2}$

21.  $\frac{24abc}{7xyz^2} \cdot \frac{14x^2yz}{9a^2}$

25.  $\frac{x + y}{3} \cdot \frac{12}{(x + y)^2}$

29.  $\frac{3b - 6}{4b + 8} \cdot \frac{5b + 10}{2 - b}$

2.  $\frac{3}{8} \cdot \frac{5}{9}$

6.  $\frac{16b}{7a} \cdot \frac{5}{4}$

10.  $\frac{7a}{12b} \cdot \frac{9b}{28}$

14.  $\frac{14y}{23x} \div 7y$

18.  $\frac{18x^2y^2}{5ab} \cdot \frac{25a^2b}{12xy}$

22.  $\frac{80x^2yz^3}{11mn^2} \cdot \frac{33mn^2}{25xyz}$

26.  $\frac{5(a - b)}{8} \cdot \frac{12}{10(a - b)}$

30.  $\frac{8y + 16}{3 - y} \cdot \frac{4y - 12}{3y + 6}$

3.  $\frac{7}{10} \div \frac{21}{25}$

7.  $\frac{14}{3a} \div \frac{7}{15a}$

11.  $\frac{9x^2}{8} \cdot \frac{4}{6x}$

15.  $6a \div \frac{24a}{35x}$

19.  $\frac{5x^2}{9y^3} \div \frac{20x}{6y}$

23.  $\frac{3ab}{8x^2} \div \frac{15b^3}{16x}$

27.  $\frac{9 - p}{7} \div \frac{4(p - 9)}{21}$

31.  $\frac{4a + 12}{a - 5} \div (a + 3)$

4.  $\frac{56}{39} \div \frac{8}{13}$

8.  $\frac{6x}{5y} \div \frac{21x}{15y}$

12.  $\frac{36p^2}{7q} \cdot \frac{14q^2}{28p^3}$

16.  $7y \div \frac{14y}{23}$

20.  $\frac{28m}{15n} \div \frac{7m^2}{3n^3}$

24.  $\frac{20mn^3}{9x^2} \div \frac{4mn}{3xy^2}$

28.  $\frac{4x - 2}{15} \div \frac{1 - 2x}{27}$

32.  $\frac{9 - 3z}{2z + 8} \div (6 - 2z)$



33.  $(x^2 - 4x + 4) \cdot \frac{18}{x^2 - 4}$
34.  $\frac{21}{a^2 - 9} \cdot (a^2 + a - 12)$
35.  $\frac{x^2 - 4}{25y} \cdot \frac{24y^2}{x + 2}$
36.  $\frac{16a^2}{b^2 - 9} \cdot \frac{b - 3}{12a^2}$
37.  $\frac{r^2 - 16}{r + 1} \div \frac{r + 4}{r^2 - 1}$
38.  $\frac{p^2 + 2p + 1}{4p - 1} \div \frac{p^2 - 1}{16p^2 - 1}$
39.  $\frac{9 - x^2}{x + y} \cdot \frac{4x + 4y}{x - 3}$
40.  $\frac{b^2 - a^2}{2a + 4b} \cdot \frac{a + b}{a - b}$
41.  $\frac{a^2 - 5a + 6}{a^2 - 9a + 20} \cdot \frac{a^2 - 5a + 4}{a^2 - 3a + 2}$
42.  $\frac{a^2 - 5a - 14}{a^2 - 9a - 36} \cdot \frac{a^2 + 10a + 21}{a^2 + 4a - 77}$
43.  $\frac{x^2 - 2x - 3}{x^2 + 3x - 4} \div \frac{x^2 - x - 6}{x^2 + x - 12}$
44.  $\frac{y^2 + 3y + 2}{y^2 + 5y + 4} \div \frac{y^2 + 5y + 6}{y^2 + 10y + 24}$
45.  $\frac{2x^2 - 15x + 7}{x^2 - 9x + 8} \cdot \frac{x^2 - 2x + 1}{x^2 - 49}$
46.  $\frac{4x^2 - 4}{3x^2 - 13x - 10} \cdot \frac{x^2 - 6x + 5}{4x + 4}$
47.  $\frac{6r^2 - r - 7}{12r^2 + 16r - 35} \div \frac{r^2 - r - 2}{2r^2 + r - 10}$
48.  $\frac{4x^2 - 9}{x^2 - 9x + 18} \div \frac{2x^2 - 5x - 12}{x^2 - 10x + 24}$
49.  $(3x^2 - 2x - 8) \div \frac{x^2 - 4}{x + 2}$
50.  $(8a^2 - 16a) \div \frac{a^3 - 16a}{a - 4}$
51.  $\frac{3x - 4}{2x + 1} \div (6x^2 - 5x - 4)$
52.  $\frac{m^2 - 3m - 10}{m^2 - 4} \div (2m^2 - 9m - 5)$
53.  $\frac{10}{a^3 - 27} \cdot \frac{a^2 + 3a - 18}{15}$
54.  $\frac{x^3 - 8}{16} \cdot \frac{24}{x^2 + 2x - 8}$
55.  $\frac{z^2 - 5z - 14}{z - 4} \div \frac{5z^3 + 40}{z^2 - z - 12}$
56.  $\frac{3b^3 + 3}{b - 2} \div \frac{b^2 + 2b + 1}{b^2 + 6b - 16}$
57.  $\frac{y^2 + 8y + 16}{y + 4} \cdot \frac{y^2 - 25}{y^2 + 9y + 20} \cdot \frac{y^2 + 5y}{y^2 - 5y}$
58.  $\frac{6m^2 - 7m + 2}{6m^2 + 5m + 1} \cdot \frac{2m^2 + m}{4m^2 - 1} \cdot \frac{12m^2 - 5m - 3}{12m^2 - 17m + 6}$

### Review exercises

Add or subtract the following. See section 1-1.

1.  $\frac{3}{4} + \frac{5}{6}$

2.  $\frac{7}{8} - \frac{5}{12}$

Completely factor the following. See sections 4-2 and 4-3.

3.  $2x^2 - 50$

4.  $x^2 + 9x - 22$

5.  $x^2 + 8x + 16$

Solve the following proportions. See section 5-4.

6.  $\frac{3}{x} = \frac{5}{8}$

7.  $\frac{5}{9} = \frac{y}{27}$

8. Write the number 0.0000789 in scientific notation. See section 3-5.

# Love The Taste. Taste The Love.

At Culver's® we can't think of anything better than serving up our creamy frozen custard and delicious classics cooked fresh the minute you order them. Which is why when we bring them to your table, they're always accompanied by a warm smile and a friendly offer to see if there's anything else we can get for you. So come on into your neighborhood Culver's and see for yourself. You might just be in love by the time you leave.





## 6-2 ■ Addition and subtraction of rational expressions

Recall that to add or subtract fractions having the same denominator, we add, or subtract, the numerators and place this sum, or difference, over the same denominator.

### Addition and subtraction properties for fractions

If  $a$ ,  $b$ , and  $c$  are real numbers,  $b \neq 0$ , then

$$\frac{a}{b} + \frac{c}{b} = \frac{a + c}{b} \quad \text{and} \quad \frac{a}{b} - \frac{c}{b} = \frac{a - c}{b}$$

### ■ Example 6-2 A

Add or subtract as indicated.

$$\begin{aligned} 1. \quad \frac{3}{11} + \frac{4}{11} &= \frac{3 + 4}{11} && \text{Add numerators} \\ &= \frac{7}{11} && 3 + 4 = 7 \end{aligned}$$

$$\begin{aligned} 2. \quad \frac{3}{7} - \frac{1}{7} &= \frac{3 - 1}{7} && \text{Subtract numerators} \\ &= \frac{2}{7} && 3 - 1 = 2 \end{aligned}$$

We use the following similar procedure to add or subtract rational expressions.

### Addition and subtraction properties for rational expressions

If  $\frac{P}{R}$  and  $\frac{Q}{R}$  are rational expressions,  $R \neq 0$ , then

$$\frac{P}{R} + \frac{Q}{R} = \frac{P + Q}{R} \quad \text{and} \quad \frac{P}{R} - \frac{Q}{R} = \frac{P - Q}{R}$$

**Note** Rational expressions having common denominators are called *like* rational expressions.

### Addition and subtraction of like rational expressions

1. Add or subtract the numerators.
2. Place the sum or difference over the common denominator.
3. Reduce the resulting rational expression to lowest terms.

### ■ Example 6-2 B

Find the indicated sum or difference. Assume all denominators are nonzero.

$$\begin{aligned} 1. \quad \frac{3}{x-2} + \frac{5}{x-2} &= \frac{3 + 5}{x-2} && \text{Add numerators and place over } x - 2 \\ &= \frac{8}{x-2} && 3 + 5 = 8 \end{aligned}$$



$$2. \frac{5y}{3y+5} - \frac{9y}{3y+5} = \frac{5y-9y}{3y+5} \\ = \frac{-4y}{3y+5}$$

Subtract numerators and place over  $3y + 5$ 

$$5y - 9y = -4y$$

$$3. \frac{2x-1}{x^2+5x+6} - \frac{4-x}{x^2+5x+6} \\ = \frac{(2x-1) - (4-x)}{x^2+5x+6} \\ = \frac{2x-1-4+x}{x^2+5x+6} \\ = \frac{3x-5}{x^2+5x+6}$$

Place numerators in parentheses and subtract

Remove parentheses and subtract

Combine like terms

**Note** Notice that when we subtracted  $4 - x$  from  $2x - 1$ , we placed parentheses around each polynomial. This step is *important* to avoid the common mistake of failing to change signs in the second expression when subtraction is involved.

$$4. \frac{2x-1}{x^2+5x+6} + \frac{4-x}{x^2+5x+6} \\ = \frac{(2x-1) + (4-x)}{x^2+5x+6} \\ = \frac{x+3}{x^2+5x+6} \\ = \frac{x+3}{(x+3)(x+2)} \\ = \frac{1}{x+2}$$

Place numerators in parentheses and add

Remove parentheses and combine like terms

Factor denominator

Reduce by common factor  $x + 3$ 

**Note** In the last step, *always* look for a possible reduction to lowest terms as we did in example 4.

► **Quick check**  $\frac{4m-5}{m^2+9} - \frac{2m-3}{m^2+9}$  ■

When one denominator is the opposite of the other, as in the indicated sum

$$\frac{2x}{3} + \frac{5}{-3},$$

where 3 and  $-3$  are opposites, we first multiply one of the expressions by  $\frac{-1}{-1}$  to obtain equivalent expressions with the same denominator.

### Example 6-2 C

Find the indicated sum or difference. Assume all denominators are not zero.

$$\begin{aligned}
 1. \quad \frac{2x}{3} + \frac{5}{-3} &= \frac{2x}{3} + \frac{-1}{-1} \cdot \frac{5}{-3} && \text{Multiply } \frac{5}{-3} \text{ by } \frac{-1}{-1} \\
 &= \frac{2x}{3} + \frac{-1(5)}{-1(-3)} && \text{Multiply numerators and denominators} \\
 &= \frac{2x}{3} + \frac{-5}{3} && \text{Common denominator of 3} \\
 &= \frac{2x + (-5)}{3} && \text{Add numerators and place over 3} \\
 &= \frac{2x - 5}{3} && \text{Definition of subtraction}
 \end{aligned}$$

$$\begin{aligned}
 2. \quad \frac{5y - 1}{y - 4} + \frac{2y + 3}{4 - y} &= \frac{5y - 1}{y - 4} + \frac{-1}{-1} \cdot \frac{2y + 3}{4 - y} && \text{Multiply } \frac{2y + 3}{4 - y} \text{ by } \frac{-1}{-1} \\
 &= \frac{5y - 1}{y - 4} + \frac{-1(2y + 3)}{-1(4 - y)} && \\
 &= \frac{5y - 1}{y - 4} + \frac{-2y - 3}{y - 4} && \text{Same denominator: } -1(4 - y) = y - 4 \\
 &= \frac{(5y - 1) + (-2y - 3)}{y - 4} && \text{Add numerators in parentheses} \\
 &= \frac{5y - 1 - 2y - 3}{y - 4} && \text{Remove parentheses} \\
 &= \frac{3y - 4}{y - 4} && \text{Combine like terms}
 \end{aligned}$$

**Note** We could have multiplied  $\frac{5y - 1}{y - 4}$  by  $\frac{-1}{-1}$ . The resulting denominator would then have been  $4 - y$  and the numerator would have been  $4 - 3y$ . We would then multiply this by  $\frac{-1}{-1}$  to obtain the same form of the answer.

$$\begin{aligned}
 3. \quad \frac{2x + 1}{x - 5} - \frac{x - 4}{5 - x} &= \frac{2x + 1}{x - 5} - \frac{-1}{-1} \cdot \frac{x - 4}{5 - x} && \text{Multiply } \frac{x - 4}{5 - x} \text{ by } \frac{-1}{-1} \\
 &= \frac{2x + 1}{x - 5} - \frac{-1(x - 4)}{-1(5 - x)} && \\
 &= \frac{2x + 1}{x - 5} - \frac{4 - x}{x - 5} && \text{Same denominator: } -1(x - 4) = 4 - x \\
 &= \frac{(2x + 1) - (4 - x)}{x - 5} && \text{Place "()" around numerators and subtract} \\
 &= \frac{2x + 1 - 4 + x}{x - 5} && \text{Definition of subtraction} \\
 &= \frac{3x - 3}{x - 5} && \text{Combine like terms}
 \end{aligned}$$

► **Quick check**  $\frac{x + 7}{x - 1} + \frac{3x + 1}{1 - x}$

**The least common denominator (LCD)**

If the fractions to be added or subtracted do not have the same denominator, we must change at least one of the fractions to an equivalent fraction so the fractions do have a common denominator. There are many such numbers we could use as a common denominator. However, the most convenient denominator to use is the smallest (least) number that is exactly divisible by each of the denominators—called the **least common denominator**, denoted by LCD. For example, the least common denominator (LCD) of the two fractions

$$\frac{5}{6} \text{ and } \frac{2}{9}$$

is 18, since 18 is the smallest (least) number that is exactly divisible by both 6 and 9.

**Finding the LCD of a set of denominators**

1. Factor each denominator completely. Write each factorization using exponential notation.
2. List each *different* factor that appears in any one of the factorizations in step 1.
3. Raise each factor of step 2 to the *greatest* power that factor has in step 1. Form the product of these factors.

**Note** The LCD of two or more rational expressions is also called the least common multiple (LCM) of the denominators.

**Example 6–2 D**

Find the LCD of rational expressions having the given denominators.

1. 6 and 9

$$\left. \begin{array}{l} 6 = 2 \cdot 3 \\ 9 = 3 \cdot 3 = 3^2 \end{array} \right\} \text{Factor each denominator}$$

The different factors are 2 and 3. The greatest power of 2 is  $2^1$  and of 3 is  $3^2$ . The LCD is  $2^1 \cdot 3^2 = 2 \cdot 9 = 18$ .

2.  $16a$  and  $8a^3$

$$\begin{aligned} 16a &= 2 \cdot 2 \cdot 2 \cdot 2 \cdot a = 2^4 \cdot a \\ 8a^3 &= 2 \cdot 2 \cdot 2 \cdot a \cdot a \cdot a = 2^3 \cdot a^3 \end{aligned}$$

Since the different factors are 2 and  $a$ , the greatest power of 2 is  $2^4$ , and the greatest power of  $a$  is  $a^3$ , the LCD is  $2^4 \cdot a^3 = 16a^3$ .

3.  $50x^3y^2$  and  $20x^2y^3$

$$\begin{aligned} 50x^3y^2 &= 2 \cdot 5 \cdot 5 \cdot x^3 \cdot y^2 = 2 \cdot 5^2 \cdot x^3 \cdot y^2 \\ 20x^2y^3 &= 2 \cdot 2 \cdot 5 \cdot x^2 \cdot y^3 = 2^2 \cdot 5 \cdot x^2 \cdot y^3 \end{aligned}$$

The different factors are 2, 5,  $x$ , and  $y$ . Since the greatest power of 2 is  $2^2$ , of 5 is  $5^2$ , of  $x$  is  $x^3$ , and of  $y$  is  $y^3$ , the LCD is  $2^2 \cdot 5^2 \cdot x^3 \cdot y^3 = 100x^3y^3$ .



4.  $x^2 + x - 12$  and  $x^2 + 2x - 8$

$$x^2 + x - 12 = (x + 4)(x - 3)$$

$$x^2 + 4x - 8 = (x + 4)(x - 2)$$

The different factors are  $x + 4$ ,  $x - 3$ , and  $x - 2$ .Each factor is carried to the first power so the LCD is  $(x + 4)(x - 3)(x - 2)$ .

5.  $x^2 - 2x + 1$ ,  $x^2 + 11x - 12$ , and  $1 - x$

$$x^2 - 2x + 1 = (x - 1)^2$$

$$x^2 + 11x - 12 = (x - 1)(x + 12)$$

$$1 - x = -1(x - 1)$$

The different factors are  $x - 1$  and  $x + 12$ . Since the greatest power of  $x - 1$  is  $(x - 1)^2$  and of  $x + 12$  is  $(x + 12)^1$ , the LCD is  $(x - 1)^2(x + 12)$ .**Note** Factors  $x - 1$  and  $1 - x$  are opposites and we will change  $1 - x$  to  $-1(x - 1)$  when working with the denominators. The factor  $-1$  is then carried along in the problem.**Quick check** Find the LCD for  $3x - 6$ ,  $x^2 - 4x + 4$ ,  $x^2 - 2x$ . ■**Mastery points****Can you**

- Add and subtract like rational expressions?
- Add and subtract rational expressions having denominators that are opposites?
- Find the least common denominator (LCD) of a set of rational expressions?

**Exercise 6-2**

Combine the given rational expressions and reduce the answer to lowest terms. Assume all denominators are nonzero. See example 6-2 B.

**Example**  $\frac{4m - 5}{m^2 + 9} - \frac{2m - 3}{m^2 + 9}$

**Solution**  $= \frac{(4m - 5) - (2m - 3)}{m^2 + 9}$

Place numerators in parentheses and subtract

$$= \frac{4m - 5 - 2m + 3}{m^2 + 9}$$

Remove parentheses and subtract in numerator

$$= \frac{2m - 2}{m^2 + 9}$$

Combine like terms in numerator

1.  $\frac{5}{x} + \frac{3}{x}$

2.  $\frac{8}{y^2} + \frac{10}{y^2}$

3.  $\frac{9}{p} - \frac{2}{p}$

4.  $\frac{18}{m^2} - \frac{5}{m^2}$

$$\boxed{5.} \frac{5x}{x+2} + \frac{9x}{x+2}$$

$$6. \frac{8y}{y-1} + \frac{-3y}{y-1}$$

$$7. \frac{x-1}{2x} - \frac{x+3}{2x}$$

$$\boxed{8.} \frac{3y-2}{y^2} - \frac{4y-1}{y^2}$$

$$9. \frac{3x+5}{x^2-1} - \frac{2x+3}{x^2-1}$$

$$10. \frac{b^2+2}{b+3} - \frac{b^2+2b-3}{b+3}$$

See example 6-2 C.

**Example**  $\frac{x+7}{x-1} + \frac{3x+1}{1-x}$

**Solution**

$$\begin{aligned} &= \frac{x+7}{x-1} + \frac{-1}{-1} \cdot \frac{3x+1}{1-x} \\ &= \frac{x+7}{x-1} + \frac{-1(3x+1)}{-1(1-x)} \\ &= \frac{x+7}{x-1} + \frac{-3x-1}{x-1} \\ &= \frac{x+7-3x-1}{x-1} \\ &= \frac{-2x+6}{x-1} \end{aligned}$$

Multiply  $\frac{3x+1}{1-x}$  by  $\frac{-1}{-1}$

$$-1(1-x) = x-1$$

Add numerators

Combine like terms

$$11. \frac{5}{7} + \frac{6}{-7}$$

$$12. \frac{9}{10} - \frac{3}{-10}$$

$$13. \frac{4}{z} - \frac{5}{-z}$$

$$14. \frac{6}{y} + \frac{9}{-y}$$

$$15. \frac{5}{x-2} + \frac{12}{2-x}$$

$$\boxed{16.} \frac{1}{x-7} - \frac{5}{7-x}$$

$$17. \frac{5y}{y-6} - \frac{4y}{6-y}$$

$$18. \frac{4z}{z-3} + \frac{z}{3-z}$$

$$19. \frac{x+1}{x-5} + \frac{2x-3}{5-x}$$

$$20. \frac{4y+3}{y-9} - \frac{2y-7}{9-y}$$

$$\boxed{21.} \frac{2y-5}{2y-3} - \frac{y+7}{3-2y}$$

$$22. \frac{z+5}{4z-3} + \frac{4z-1}{3-4z}$$

$$23. \frac{2x+5}{5-2x} + \frac{x+9}{2x-5}$$

$$24. \frac{5-y}{6-5y} - \frac{9y+1}{5y-6}$$

Find the least common denominator (LCD) of rational expressions having the following denominators.  
See example 6-2 D.

**Example**  $3x-6$ ,  $x^2-4x+4$ , and  $x^2-2x$

**Solution**

$$\begin{aligned} 3x-6 &= 3(x-2) \\ x^2-4x+4 &= (x-2)^2 \\ x^2-2x &= x(x-2) \end{aligned} \quad \left\{ \begin{array}{l} \text{Factor each denominator} \end{array} \right.$$

LCD is  $3x(x-2)^2$

$$25. 6x \text{ and } 9x$$

$$26. 8a \text{ and } 12a$$

$$27. 16x^2 \text{ and } 24x$$

$$28. 6b^2 \text{ and } 14b$$

$$29. 28y^2 \text{ and } 35y^3$$

$$30. 9z^3 \text{ and } 7z^4$$

$$31. 32a^2 \text{ and } 64a^4$$

$$\boxed{32.} 4x^2, 3x, \text{ and } 8x^3$$

$$33. 10a^2, 12a^3, \text{ and } 9a$$

$$34. 4x-2 \text{ and } 2x-1$$

$$35. x-4 \text{ and } 3x-12$$

$$36. 6x-12 \text{ and } 9x-18$$

37.  $18y^3$  and  $9y - 36$       38.  $32z^2$  and  $16z - 32$       39.  $a^2 + a$  and  $a^2 - 1$   
 40.  $(z - 1)^2$  and  $z^2 - 1$       41.  $8a + 16$  and  $a^2 + 3a + 2$       42.  $9p - 18$  and  $p^2 - 7p + 10$   
 43.  $a^2 - 5a + 6$  and  $a^2 - 4$       44.  $y^2 - y - 12$  and  $y^2 + 6y + 9$   
 45.  $a^2 - 9$  and  $a^2 - 5a + 6$       46.  $p^2 - 9$ ,  $p^2 + p - 6$ , and  $p^2 - 4p + 4$   
 47.  $x^2 - 49$ ,  $7 - x$ , and  $2x + 14$       48.  $5 - y$ ,  $y^2 - 25$ , and  $y^2 - 10y + 25$

### Review exercises

1. The statement  $4(y + 3) = 4(3 + y)$  demonstrates what property of real numbers? See section 1-8.

Factor the following expressions. See sections 4-2, 4-3, and 4-4.

2.  $5y^2 - 20$       3.  $x^2 + 20x + 100$       4.  $3y^2 - y - 4$

Find the solution set of the following equations. See sections 2-6 and 4-7.

5.  $4(x + 3) = 5(4 - 3x)$       6.  $x^2 - 2x = 15$

## 6-3 ■ Addition and subtraction of rational expressions

Now that we can find the least common denominator (LCD) of a group of rational expressions, let us review the process for changing a fraction (or rational expression) to an equivalent fraction with a new denominator.

### ■ Example 6-3 A

1. Change  $\frac{7}{15}$  to an equivalent fraction having denominator 60.

We want  $\frac{7}{15} = \frac{?}{60}$ .

Since  $60 = 15 \cdot 4$  (factor 4 is found by dividing  $60 \div 15 = 4$ ), we multiply the given fraction by  $\frac{4}{4}$  ( $\frac{4}{4} = 1$ ).

$$\begin{aligned} \frac{7}{15} &= \frac{7}{15} \cdot \frac{4}{4} \\ &= \frac{7 \cdot 4}{15 \cdot 4} && \begin{array}{l} \text{Multiply numerators} \\ \text{Multiply denominators} \end{array} \\ &= \frac{28}{60} \end{aligned}$$

$$\text{Thus, } \frac{7}{15} = \frac{28}{60}.$$



2. Change  $\frac{x+1}{x-4}$  to an equivalent rational expression having denominator  $x^2 - 2x - 8$ .

$$\text{We want } \frac{x+1}{x-4} = \frac{?}{x^2 - 2x - 8}.$$

Since  $x^2 - 2x - 8 = (x-4)(x+2)$ , we multiply the given rational expression by  $\frac{x+2}{x+2}$ .

$$\begin{aligned} \frac{x+1}{x-4} &= \frac{x+1}{x-4} \cdot \frac{x+2}{x+2} \\ &= \frac{(x+1)(x+2)}{(x-4)(x+2)} && \text{Multiply numerators} \\ &= \frac{x^2 + 3x + 2}{x^2 - 2x - 8} && \text{Multiply denominators} \\ &= \frac{x^2 + 3x + 2}{x^2 - 2x - 8} && \text{Perform indicated operations} \end{aligned}$$

$$\text{Thus, } \frac{x+1}{x-4} = \frac{x^2 + 3x + 2}{x^2 - 2x - 8}.$$

Once equivalent rational expressions are obtained with the LCD as the denominator, we add or subtract as previously learned. Use the following steps to add or subtract rational expressions having unlike denominators.

### Addition and subtraction of rational expressions having different denominators

1. Find the LCD of the rational expressions.
2. Write each rational expression as an equivalent rational expression with the LCD as the denominator.
3. Perform the indicated addition or subtraction as before.
4. Reduce the results to lowest terms.

### Example 6-3 B

Add the following rational expressions. Assume the denominators are not equal to zero. Reduce all answers to lowest terms.

$$1. \frac{5x}{8} + \frac{7x}{12}$$

$$\left. \begin{array}{l} 8 = 2 \cdot 2 \cdot 2 \\ 12 = 2 \cdot 2 \cdot 3 \end{array} \right\} \text{LCD} = 2 \cdot 2 \cdot 2 \cdot 3 = 24 \quad \text{Find the LCD}$$

$$\text{Since } \frac{24}{8} = 3 \text{ and } \frac{24}{12} = 2$$

$$\begin{aligned} \frac{5x}{8} + \frac{7x}{12} &= \frac{5x}{8} \cdot \frac{3}{3} + \frac{7x}{12} \cdot \frac{2}{2} && \text{Multiply } \frac{5x}{8} \text{ by } \frac{3}{3} \text{ and } \frac{7x}{12} \text{ by } \frac{2}{2} \\ &= \frac{15x}{24} + \frac{14x}{24} && \text{Multiply numerators and denominators} \\ &= \frac{15x + 14x}{24} && \text{Add numerators} \\ &= \frac{29x}{24} && \text{Combine as indicated} \end{aligned}$$

$$2. \frac{15}{4x^2} + \frac{25}{18x}$$

$$\left. \begin{array}{l} 4x^2 = 2 \cdot 2 \cdot x^2 \\ 18x = 2 \cdot 3 \cdot 3 \cdot x \end{array} \right\} \text{LCD} = 2 \cdot 2 \cdot 3 \cdot 3 \cdot x^2 = 36x^2 \quad \text{Find the LCD}$$

$$\text{Since } \frac{36x^2}{4x^2} = 9 \text{ and } \frac{36x^2}{18x} = 2x$$

$$\frac{15}{4x^2} + \frac{25}{18x} = \frac{15}{4x^2} \cdot \frac{9}{9} + \frac{25}{18x} \cdot \frac{2x}{2x}$$

Multiply  $\frac{15}{4x^2}$  by  $\frac{9}{9}$  and  $\frac{25}{18x}$  by  $\frac{2x}{2x}$

$$= \frac{135}{36x^2} + \frac{50x}{36x^2}$$

Multiply numerators and denominators

$$= \frac{135 + 50x}{36x^2}$$

Add numerators

$$3. \frac{3y+2}{y^2-16} + \frac{y-4}{3y+12}$$

$$\left. \begin{array}{l} y^2 - 16 = (y+4)(y-4) \\ 3y + 12 = 3(y+4) \end{array} \right\} \text{LCD} = 3(y+4)(y-4) \quad \text{Find the LCD}$$

$$\text{Since } \frac{3(y+4)(y-4)}{(y+4)(y-4)} = 3 \text{ and } \frac{3(y+4)(y-4)}{3(y+4)} = y-4, \text{ then}$$

$$\frac{3y+2}{y^2-16} + \frac{y-4}{3y+12}$$

$$= \frac{3y+2}{(y+4)(y-4)} \cdot \frac{3}{3} + \frac{y-4}{3(y+4)} \cdot \frac{y-4}{y-4}$$

Multiply  $\frac{3y+2}{(y+4)(y-4)}$   
by  $\frac{3}{3}$  and  $\frac{y-4}{3(y+4)}$  by  
 $\frac{y-4}{y-4}$

$$= \frac{3(3y+2)}{3(y+4)(y-4)} + \frac{(y-4)(y-4)}{3(y+4)(y-4)}$$

Multiply numerators  
and denominators

$$= \frac{(9y+6) + (y^2-8y+16)}{3(y+4)(y-4)}$$

Add numerators

$$= \frac{y^2 + y + 22}{3(y+4)(y-4)}$$

Remove parentheses  
and combine

**Note** When the numerators have two or more terms as in example 3, we place the quantities in parentheses when we add the numerators. This is a good practice to avoid a *most common* mistake when subtracting.

► **Quick check** Add the rational expressions  $\frac{20}{3y} + \frac{25}{12y^2}$ .



### Example 6-3 C

Subtract the following rational expressions. Assume the denominators are not equal to zero. Reduce to lowest terms.

$$1. \frac{5y}{2y-1} - \frac{y+1}{y+2}$$

The LCD of the rational expressions is  $(2y-1)(y+2)$ .

$$\begin{aligned} \frac{5y}{2y-1} - \frac{y+1}{y+2} &= \frac{5y}{2y-1} \cdot \frac{y+2}{y+2} - \frac{y+1}{y+2} \cdot \frac{2y-1}{2y-1} \\ &= \frac{5y(y+2)}{(2y-1)(y+2)} - \frac{(y+1)(2y-1)}{(2y-1)(y+2)} \\ &= \frac{(5y^2 + 10y) - (2y^2 + y - 1)}{(2y-1)(y+2)} \\ &= \frac{5y^2 + 10y - 2y^2 - y + 1}{(2y-1)(y+2)} \\ &= \frac{3y^2 + 9y + 1}{(2y-1)(y+2)} \end{aligned}$$

Multiply  $\frac{5y}{2y-1}$  by  $\frac{y+2}{y+2}$  and  $\frac{y+1}{y+2}$  by  $\frac{2y-1}{2y-1}$

Multiply numerators and denominators

Subtract numerators

Remove parentheses and change signs when subtracting

Combine like terms

Don't forget the parentheses

**Note** The numerator  $3y^2 + 9y + 1$  cannot be factored so we are unable to reduce. We should always check this!

$$2. \frac{5x-4}{x^2-2x+1} - \frac{3x}{x^2+4x-5}$$

$$\left. \begin{aligned} x^2 - 2x + 1 &= (x-1)^2 \\ x^2 + 4x - 5 &= (x+5)(x-1) \end{aligned} \right\} \text{LCD} = (x-1)^2(x+5) \quad \text{Find the LCD}$$

$$\begin{aligned} \frac{5x-4}{x^2-2x+1} - \frac{3x}{x^2+4x-5} &= \frac{5x-4}{(x-1)^2} - \frac{3x}{(x+5)(x-1)} \\ &= \frac{5x-4}{(x-1)^2} \cdot \frac{x+5}{x+5} - \frac{3x}{(x+5)(x-1)} \cdot \frac{x-1}{x-1} \end{aligned}$$

Factor denominators

Multiply  $\frac{5x-4}{(x-1)^2}$  by  $\frac{x+5}{x+5}$  and

$\frac{3x}{(x+5)(x-1)}$  by  $\frac{x-1}{x-1}$

$\frac{3x}{(x+5)(x-1)}$  by  $\frac{x-1}{x-1}$

$$\begin{aligned} &= \frac{(5x-4)(x+5)}{(x-1)^2(x+5)} - \frac{3x(x-1)}{(x-1)^2(x+5)} \\ &= \frac{(5x^2 + 21x - 20) - (3x^2 - 3x)}{(x-1)^2(x+5)} \\ &= \frac{5x^2 + 21x - 20 - 3x^2 + 3x}{(x-1)^2(x+5)} \\ &= \frac{2x^2 + 24x - 20}{(x-1)^2(x+5)} \end{aligned}$$

Multiply numerators and denominators

Subtract numerators

Remove parentheses and change signs

Combine like terms

Don't forget the parentheses



► **Quick check** Subtract  $\frac{y+1}{y^2-y-12} - \frac{3y+2}{y^2-9y+20}$

### Problem solving

#### ■ Example 6-3 D

Set up a rational expression for the following word statements.

1. If Dick can mow his lawn in  $h$  hours, what part of the lawn can he mow in 2 hours?

We must find what part of the total time,  $h$  hours, is 2 hours. Thus, Dick can mow the fractional part

$$\frac{2}{h}$$

of the lawn in 2 hours.

2. If the area ( $A$ ) of a rectangle is  $n$  square inches, what is the expression for the length,  $\ell$ , if the width,  $w$ , is 12 inches? ( $A = \ell \cdot w$ )

Using  $A = \ell \cdot w$ , we have  $\ell = \frac{A}{w}$  and so the length

$$\ell = \frac{n}{12} \text{ inches.} \quad \text{Replace } A \text{ with } n \text{ and } w \text{ with } 12$$

► **Quick check** If the area of a rectangle is  $p$  square yards, write an expression for the width,  $w$ , if the length is 5 yards.

### Mastery points

Can you

- Add and subtract rational expressions having unlike denominators?

### Exercise 6-3

Perform the indicated addition and reduce the answer to lowest terms. Assume all denominators are not equal to zero. See example 6-3 B.

**Example**  $\frac{20}{3y} + \frac{25}{12y^2}$

**Solution**  $\left. \begin{array}{l} 3y = 3 \cdot y \\ 12y^2 = 3 \cdot 2 \cdot 2 \cdot y \cdot y \end{array} \right\} \text{LCD} = 3 \cdot 2 \cdot 2 \cdot y \cdot y = 12y^2$

$\frac{12y^2}{3y} = 4y$  and the denominator of the second expression is the LCD,  $12y^2$ .

$$\begin{aligned} \frac{20}{3y} + \frac{25}{12y^2} &= \frac{20}{3y} \cdot \frac{4y}{4y} + \frac{25}{12y^2} \\ &= \frac{80y}{12y^2} + \frac{25}{12y^2} \\ &= \frac{80y + 25}{12y^2} \end{aligned}$$

Multiply numerator and denominator of  $\frac{20}{3y}$  by  $4y$

Perform indicated operations

Add numerators and place over common denominator

1.  $\frac{x}{6} + \frac{3}{4}$
2.  $\frac{3z}{10} + \frac{2z}{15}$
3.  $\frac{2x-1}{16} + \frac{x+2}{24}$
4.  $\frac{4}{3x} + \frac{5}{2x}$
5.  $\frac{3a+1}{a} + \frac{2a-3}{3a}$
6.  $\frac{4}{x-1} + \frac{5}{x+3}$
7.  $\frac{8}{y+4} + \frac{7}{y-5}$
8.  $\frac{x}{x+2} + \frac{3x}{4x-1}$
9.  $\frac{15}{5y-10} + \frac{14}{2y-4}$
10.  $\frac{21}{6x+12} + \frac{15}{2x+4}$
11.  $\frac{12}{x^2-4} + \frac{7}{4x-8}$
12.  $\frac{16}{2y+6} + \frac{5}{y^2-9}$
13.  $5 + \frac{4x}{x+8}$
14.  $9 + \frac{y+9}{y-1}$
15.  $\frac{x}{x-1} + \frac{3x}{x^2-1}$
16.  $\frac{2y}{y^2-16} + \frac{5y}{2y-8}$
17.  $\frac{4}{x^2-x-6} + \frac{5}{x^2-9}$
18.  $\frac{6}{x^2-4x-12} + \frac{5}{x^2-36}$
19.  $\frac{2y}{y^2-6y+9} + \frac{5y}{y^2-2y-3}$
20.  $\frac{4z}{z^2+z-20} + \frac{z}{z^2-8z+16}$
21.  $\frac{y-2}{y^2-3y-10} + \frac{y+1}{y^2-y-6}$
22.  $\frac{2x+1}{x^2+6x+5} + \frac{4x-3}{x^2-x-30}$

Perform the indicated subtraction and reduce to lowest terms. Assume all denominators are not zero. See example 6-3 C.

**Example**  $\frac{y+1}{y^2-y-12} - \frac{3y+2}{y^2-9y+20}$

**Solution**  $\left. \begin{array}{l} y^2-y-12 = (y-4)(y+3) \\ y^2-9y+20 = (y-4)(y-5) \end{array} \right\} \text{LCD is } (y-4)(y+3)(y-5).$

$$\begin{aligned}
 & \frac{y+1}{y^2-y-12} - \frac{3y+2}{y^2-9y+20} \\
 &= \frac{y+1}{(y-4)(y+3)} \cdot \frac{y-5}{y-5} - \frac{3y+2}{(y-4)(y-5)} \cdot \frac{y+3}{y+3} \\
 &= \frac{(y+1)(y-5)}{(y-4)(y+3)(y-5)} - \frac{(3y+2)(y+3)}{(y-4)(y+3)(y-5)} \\
 &= \frac{(y^2-4y-5) - (3y^2+11y+6)}{(y-4)(y+3)(y-5)} \\
 &= \frac{y^2-4y-5-3y^2-11y-6}{(y-4)(y+3)(y-5)} \\
 &= \frac{-2y^2-15y-11}{(y-4)(y+3)(y-5)}
 \end{aligned}$$

Multiply numerator and denominator of  $\frac{y+1}{(y-4)(y+3)}$  by  $y-5$  and of  $\frac{3y+2}{(y-4)(y-5)}$  by  $y+3$

Multiply numerators and denominators

Multiply in each numerator, place parentheses around each product, and subtract

Remove parentheses, change signs, and subtract

Combine like terms in numerator

Don't forget the parentheses

**Note** The denominator is left in factored form in case the answer can be reduced.

23.  $\frac{y}{9} - \frac{5}{6}$

26.  $\frac{7}{12z} - \frac{10}{9z}$

29.  $\frac{2x+5}{6x} - \frac{x-5}{9x}$

32.  $\frac{7}{4x-6} - \frac{12}{3x+9}$

35.  $9 - \frac{6}{x+8}$

38.  $\frac{4y}{5y-4} - 10$

41.  $\frac{20}{y^2-2y-24} - \frac{8}{y^2+y-12}$

43.  $\frac{2a-3}{a^2-5a+6} - \frac{3a}{a-2}$

Add and subtract as indicated.

44.  $\frac{13}{12b} - \frac{2}{9b} + \frac{5}{4b}$

47.  $\frac{4a}{5} + \frac{7a}{15} - \frac{a}{9}$

50.  $\frac{4a}{a^2+2a-15} + \frac{3a}{2a^2+11a+5} - \frac{5a}{2a^2-5a-3}$

24.  $\frac{5y}{12} - \frac{y}{8}$

27.  $\frac{5a+3}{12} - \frac{a-4}{10}$

30.  $\frac{4y-1}{3y} - \frac{2y-3}{15y}$

33.  $\frac{12}{3y+6} - \frac{11}{7y+14}$

36.  $12 - \frac{7}{z-12}$

39.  $\frac{-3}{a^2-5a+6} - \frac{3}{a^2-4}$

42.  $\frac{2p}{p^2-9p+20} - \frac{5p-2}{p-5}$

25.  $\frac{9}{14y} - \frac{1}{21y}$

28.  $\frac{2x+9}{8} - \frac{x-7}{20}$

31.  $\frac{7}{2x-3} - \frac{6}{x-5}$

34.  $\frac{14}{5x-15} - \frac{8}{2x-6}$

37.  $\frac{2x}{3x+1} - 9$

40.  $\frac{8}{x^2-25} - \frac{7}{x^2+3x-10}$

45.  $\frac{5}{6z} + \frac{4}{8z} - \frac{1}{4z}$

48.  $\frac{a-1}{5} - \frac{2a+3}{15} + \frac{3a-1}{25}$

46.  $\frac{3x}{8} - \frac{2x}{5} + \frac{7x}{10}$

49.  $\frac{5b+1}{6} + \frac{3b-2}{9} - \frac{b+1}{12}$

51.  $\frac{5}{z^2-4} - \frac{z}{z^2-1} + \frac{4}{z^2+z-2}$

Solve the following word problems.

52. For a lens maker to manufacture lenses that will refract light by exactly the right amounts, the following formula is used:

$$\frac{1}{f} = (u-1) \left( \frac{1}{R_1} + \frac{1}{R_2} \right)$$

Add the expressions containing  $R_1$  and  $R_2$ .

53. Women A, B, and C can complete a given job in
- $a$
- ,
- $b$
- , and
- $c$
- hours, respectively. Working together they can complete in one hour
- $\frac{1}{a} + \frac{1}{b} + \frac{1}{c}$
- of the job.

By combining, obtain a single expression for what they can do together in one hour.

54. In electricity, the total resistance of any parallel circuit may be given by

$$\frac{1}{R_t} = \frac{I_1}{E_1} + \frac{I_2}{E_2} + \frac{I_3}{E_3}$$

Combine the expression in the right member.



See example 6–3 D.

**Example** If the area of a rectangle is  $p$  square yards, write an expression for the width,  $w$ , if the length is 5 yards.

**Solution** Using  $A = \ell \cdot w$ , we are given that  $A = p$  square yards and  $\ell = 5$  yards. Substituting, we obtain

$$p = 5 \cdot w$$

and solving for  $w$ , we divide each member by 5. Thus  $w = \frac{p}{5}$ .

55. A faucet when fully open, can fill the sink in  $m$  minutes. What part of the sink can it fill in 3 minutes?
56. An inlet pipe to a swimming pool can fill the pool in  $h$  hours. What part of the pool can it fill in 9 hours?
57. An outlet pipe can drain a swimming pool in 36 hours. What part of the pool can it drain in  $h$  hours?
58. Jane can paint her house in  $h$  hours. What part of the house can she paint in 1 hour?
59. The product of two numbers is 48. If one of the numbers is  $m$ , what is the other number?
60. The area of a rectangle is 54 square centimeters. What is the length  $\ell$  if the width is  $w$  centimeters?
61. The area of a rectangle is  $A$  square feet. What is the width if the length is 23 feet?
62. The area of a triangle is 21 square yards. If the triangle has a base length  $b$ , what is the altitude  $h$  of the triangle?  $\left(A = \frac{1}{2}bh\right)$
63. The area of a triangle is  $A$  square rods. If the altitude  $h$  is 9 rods, what is the length of the base  $b$ ? (See problem 62.)
64. John drives 25 miles in  $h$  hours. At what speed,  $r$ , did he travel? [Hint: Use distance traveled ( $d$ ) = rate ( $r$ )  $\times$  time ( $t$ ).]
65. Mabel travels  $d$  miles at a rate of 55 miles per hour. Write an expression for the time  $t$  that she traveled. (See problem 64.)
66. What is the reciprocal of the natural number  $n$ ?
67. What is the reciprocal of the fraction  $\frac{a}{b}$ ?

### Review exercises

Completely factor the following expressions. See sections 4–2, 4–3, and 4–4.

1.  $x^2 - 14x + 49$

2.  $2x^2 - 11x + 5$

3.  $4x^2 - 16$

Multiply the following expressions. See section 3–2.

4.  $(x + 9)(x - 9)$

5.  $(4x + 3)^2$

6.  $(2x + 1)(x - 8)$

Find the LCD of expressions having the following denominators. See section 6–2.

7. 16, 12, 6


8.  $4x$ ,  $2x^2$ , 6

9.  $x^2 - 9$ ;  $(x - 3)^2$ ;  $x + 3$

Add or subtract the following. See section 6–3.

10.  $\frac{5}{x} + \frac{3}{2x}$

11.  $\frac{5}{x - 2} - \frac{3}{x - 1}$



# Extra Credit Rocks

Sign up for a Discover® Student Card today and enjoy:

- 0% Intro APR\* on Purchases for 6 Months
- No Annual Fee
- Easiest Online Account Management Options
- Full 5% *Cashback Bonus*®\* on Get More purchases in popular categories all year
- Up to 1% *Cashback Bonus*®\* on all your other purchases
- Unlimited cash rewards that never expire as long as you use your Card

**APPLY NOW**

**DISCOVER**  
CARD

\*View Discover® Card Rates, Fees, Rewards and Other Important Information.



## 6-4 ■ Complex fractions

**A complex fraction**

A **complex fraction** is a fraction (rational expression) whose numerator, denominator, or both contain fractions (or rational expressions). The expressions

$$\frac{\frac{3}{4}}{\frac{5}{6}}, \quad \frac{\frac{1}{3} + 2}{1 - \frac{1}{2}}, \quad \text{and} \quad \frac{\frac{x-1}{x-2}}{\frac{x+3}{x}}$$

are examples of complex fractions. Given a complex fraction, we simplify the fraction by *eliminating the fractions within the numerator and/or the denominator* to obtain a simple fraction.

We name the parts of a complex fraction as shown in the following examples.

$$\begin{array}{ll} \left. \begin{array}{l} \frac{3}{x} + \frac{4}{y} \end{array} \right\} & \text{Primary numerator} \qquad \begin{array}{l} \frac{3}{x} + \frac{4}{y} \leftarrow \text{Secondary numerators} \\ \qquad \qquad \qquad \leftarrow \text{Secondary denominators} \end{array} \\ \left. \begin{array}{l} \frac{1}{x} - \frac{2}{y} \end{array} \right\} & \text{Primary denominator} \qquad \begin{array}{l} \frac{1}{x} - \frac{2}{y} \leftarrow \text{Secondary numerators} \\ \qquad \qquad \qquad \leftarrow \text{Secondary denominators} \end{array} \end{array}$$

To simplify a complex fraction, we use one of the following methods.

**Simplifying a complex fraction**

**Method 1** Multiply the primary numerator and the primary denominator by the LCD of the secondary denominators and reduce if possible.

**Method 2** Form a single fraction in the numerator and in the denominator and divide the primary numerator by the primary denominator.

■ **Example 6-4 A**

Simplify each complex fraction.

$$1. \quad \frac{\frac{3}{4}}{\frac{5}{6}}$$

*Method 1*

$$\begin{aligned} \frac{\frac{3}{4}}{\frac{5}{6}} &= \frac{\frac{3}{4} \cdot 12}{\frac{5}{6} \cdot 12} \\ &= \frac{3 \cdot \frac{12}{4}}{5 \cdot \frac{12}{6}} \\ &= \frac{3 \cdot 3}{5 \cdot 2} \\ &= \frac{9}{10} \end{aligned}$$

Multiply primary numerator and primary denominator by the LCD of secondary denominators 4 and 6, which is 12

Divide 4 into 12

Divide 6 into 12

Multiply in numerator and denominator



*Method 2*

$$\begin{aligned}
 \frac{\frac{3}{4}}{\frac{5}{6}} &= \frac{3}{4} \div \frac{5}{6} = \frac{3}{4} \cdot \frac{6}{5} \\
 &= \frac{3 \cdot 6}{4 \cdot 5} \\
 &= \frac{18}{20} \\
 &= \frac{9}{10}
 \end{aligned}$$

Multiply by the reciprocal of  $\frac{5}{6}$ 

Multiply numerators and denominators as indicated

Multiply and reduce to lowest terms

Reduce to lowest terms

$$2. \frac{\frac{3}{a} - 1}{1 + \frac{4}{b}}$$

*Method 1*

The LCD of the secondary denominators,  $a$  and  $b$ , is  $ab$ . We multiply the primary numerator,  $\frac{3}{a} - 1$ , and the primary denominator,  $1 + \frac{4}{b}$ , by  $ab$ .

Remember to multiply the LCD times *every term* in the numerator and the denominator.

$$\begin{aligned}
 \frac{\frac{3}{a} - 1}{1 + \frac{4}{b}} &= \frac{\left(\frac{3}{a} - 1\right) \cdot ab}{\left(1 + \frac{4}{b}\right) \cdot ab} \\
 &= \frac{\frac{3}{a} \cdot ab - 1 \cdot ab}{1 \cdot ab + \frac{4}{b} \cdot ab} \\
 &= \frac{3b - ab}{ab + 4a} \\
 &= \frac{b(3 - a)}{a(b + 4)}
 \end{aligned}$$

Multiply each term of numerator and denominator by  $ab$ 

Perform indicated multiplications

Factor numerator and denominator

*Method 2*

We first change the primary numerator,  $\frac{3}{a} - 1$ , and the primary denominator,  $1 + \frac{4}{b}$ , to single fractions. Thus,

$$\frac{3}{a} - 1 = \frac{3}{a} - \frac{a}{a} = \frac{3 - a}{a}$$

Combine in primary numerator

and

$$1 + \frac{4}{b} = \frac{b}{b} + \frac{4}{b} = \frac{b + 4}{b}$$

Combine in primary denominator

$$\begin{aligned}\frac{\frac{3}{a} - 1}{1 + \frac{4}{b}} &= \frac{\frac{3-a}{a}}{\frac{b+4}{b}} = \frac{3-a}{a} \cdot \frac{b}{b+4} && \text{Multiply by the reciprocal of } \frac{b+4}{b} \\ &= \frac{b(3-a)}{a(b+4)}\end{aligned}$$

**Note** Always form a single fraction in the numerator and in the denominator before we invert and multiply. That is,

$$\frac{\frac{3}{a} - 1}{1 + \frac{4}{b}} \neq \left(\frac{3}{a} - 1\right) \cdot \left(1 + \frac{b}{4}\right)$$

As you can see, both methods will work. We will use only Method 1 in the remaining examples. Choice of method is left up to the student.

$$3. \quad \frac{\frac{4}{x} - \frac{3}{y}}{\frac{5}{x} + \frac{7}{y}}$$

The LCD of the secondary denominators,  $x$  and  $y$ , is  $xy$ . We multiply the primary numerator,  $\frac{4}{x} - \frac{3}{y}$ , and the primary denominator,  $\frac{5}{x} + \frac{7}{y}$ , by  $xy$  and get

$$\begin{aligned}\frac{\frac{4}{x} - \frac{3}{y}}{\frac{5}{x} + \frac{7}{y}} &= \frac{\left(\frac{4}{x} - \frac{3}{y}\right) \cdot xy}{\left(\frac{5}{x} + \frac{7}{y}\right) \cdot xy} = \frac{\frac{4}{x} \cdot xy - \frac{3}{y} \cdot xy}{\frac{5}{x} \cdot xy + \frac{7}{y} \cdot xy} && \text{Multiply each term of} \\ &&& \text{numerator and} \\ &&& \text{denominator by } xy \\ &= \frac{4y - 3x}{5y + 7x} && \text{Perform indicated} \\ &&& \text{multiplications}\end{aligned}$$

$$4. \quad \frac{\frac{a^2 - x^2}{x}}{\frac{a + x}{x^2}}$$

The LCD of the secondary denominators,  $x$  and  $x^2$ , is  $x^2$ . We then multiply the primary numerator,  $\frac{a^2 - x^2}{x}$ , and the primary denominator,  $\frac{a + x}{x^2}$ , by  $x^2$  and get

$$\begin{aligned}\frac{\frac{a^2 - x^2}{x}}{\frac{a + x}{x^2}} &= \frac{\left(\frac{a^2 - x^2}{x}\right) \cdot x^2}{\left(\frac{a + x}{x^2}\right) \cdot x^2} = \frac{(a^2 - x^2) \cdot x}{a + x} && \text{Factor the numerator} \\ &= \frac{(a + x)(a - x) \cdot x}{a + x} && \text{Reduce by the} \\ &= (a - x) \cdot x && \text{common factor} \\ &= ax - x^2 && \text{Multiply as indicated}\end{aligned}$$

**Note** In example 4, we proceeded to *reduce before* performing the multiplication in the numerator. *Always do this, if possible.*

► **Quick check** Simplify  $\frac{\frac{3}{4} + \frac{2}{3}}{\frac{5}{6} - \frac{1}{12}}$

### Mastery points

Can you

- Simplify complex fractions?

## Exercise 6-4

Simplify each complex fraction to a simple fraction. See example 6-4 A.

**Example**

$$\frac{\frac{3}{4} + \frac{2}{3}}{\frac{5}{6} - \frac{1}{12}}$$

**Solution** Method 1

$$\begin{aligned} \frac{\frac{3}{4} + \frac{2}{3}}{\frac{5}{6} - \frac{1}{12}} &= \frac{\left(\frac{3}{4} + \frac{2}{3}\right)12}{\left(\frac{5}{6} - \frac{1}{12}\right)12} = \frac{\frac{3}{4} \cdot 12 + \frac{2}{3} \cdot 12}{\frac{5}{6} \cdot 12 - \frac{1}{12} \cdot 12} \\ &= \frac{9 + 8}{10 - 1} \\ &= \frac{17}{9} \end{aligned}$$

Multiply numerator and denominator by 12

Multiply as indicated

Combine in the numerator and the denominator

Method 2

$$\begin{aligned} \frac{\frac{3}{4} + \frac{2}{3}}{\frac{5}{6} - \frac{1}{12}} &= \frac{\frac{9}{12} + \frac{8}{12}}{\frac{10}{12} - \frac{1}{12}} = \frac{\frac{17}{12}}{\frac{9}{12}} \\ &= \frac{17}{12} \div \frac{9}{12} \\ &= \frac{17}{12} \cdot \frac{12}{9} \\ &= \frac{17}{9} \end{aligned}$$

Add in the numerator

Subtract in the denominator

Multiply by the reciprocal of  $\frac{9}{12}$

1.  $\frac{\frac{2}{3}}{\frac{4}{5}}$

2.  $\frac{\frac{7}{8}}{\frac{5}{6}}$

3.  $\frac{\frac{4}{3}}{\frac{8}{9}}$

4.  $\frac{\frac{9}{10}}{\frac{7}{6}}$

5.  $\frac{1 + \frac{3}{5}}{2 - \frac{1}{5}}$



$$6. \frac{5 - \frac{3}{4}}{1 + \frac{5}{8}}$$

$$7. \frac{\frac{7}{2} + \frac{4}{5}}{\frac{3}{14} - \frac{5}{7}}$$

$$8. \frac{10}{4 - \frac{11}{12}}$$

$$9. \frac{4 + \frac{3}{5}}{6}$$

$$10. \frac{10 - \frac{7}{8}}{3}$$

$$11. \frac{\frac{3}{4} + \frac{5}{8}}{\frac{1}{2} - \frac{1}{4}}$$

$$12. \frac{\frac{6}{7} - \frac{5}{14}}{\frac{3}{14} - \frac{5}{7}}$$

$$13. \frac{x + \frac{1}{4}}{x - \frac{3}{4}}$$

$$14. \frac{y - \frac{5}{6}}{y + \frac{1}{2}}$$

$$15. \frac{\frac{1}{a} + 3}{\frac{2}{a} - 4}$$

$$16. \frac{5 - \frac{3}{b}}{6 + \frac{5}{b}}$$

$$17. \frac{\frac{3}{a^2} + 4}{5 - \frac{3}{a}}$$

$$18. \frac{\frac{5}{x} - 5}{6 + \frac{4}{x^3}}$$

$$19. \frac{a - \frac{3}{b}}{a + \frac{4}{b}}$$

$$20. \frac{x + \frac{4}{y}}{x - \frac{5}{y}}$$

$$21. \frac{\frac{1}{x} + \frac{1}{y}}{\frac{1}{x} - \frac{1}{y}}$$

$$22. \frac{\frac{3}{x^2} - \frac{4}{y}}{\frac{5}{x} + \frac{2}{y^2}}$$

$$23. \frac{x + y}{\frac{1}{x} + \frac{1}{y}}$$

$$24. \frac{a - b}{\frac{2}{a} + \frac{3}{b}}$$

$$25. \frac{\frac{4}{a^2} - \frac{5}{b}}{a - b}$$

$$26. \frac{\frac{6}{x^2} - \frac{5}{y^2}}{x + y}$$

$$27. \frac{\frac{1}{x+y} - \frac{1}{x-y}}{\frac{1}{x+y} + \frac{1}{x-y}}$$

$$28. \frac{\frac{a}{a+b} - \frac{b}{a-b}}{\frac{b}{a+b} + \frac{a}{a-b}}$$

$$29. \frac{\frac{x+y}{x-y} + \frac{x-y}{x+y}}{x^2 - y^2}$$

$$30. \frac{\frac{b}{a+b} - \frac{a}{a-b}}{a^2 - b^2}$$

$$31. \frac{\frac{2}{ab} + \frac{3}{ab^2}}{a^2b^2}$$

$$32. \frac{\frac{5}{x^2y^2} - \frac{4}{xy}}{xy}$$

$$33. \frac{\frac{2}{a+3} + \frac{1}{a-2}}{\frac{3}{a-2} - \frac{4}{a+3}}$$

$$34. \frac{\frac{7}{x-4} - \frac{5}{x+3}}{\frac{5}{x+3} + \frac{9}{x-4}}$$

$$35. \frac{\frac{5}{x^2 - x - 12}}{\frac{4}{x+3} - \frac{5}{x-4}}$$

$$36. \frac{\frac{7}{b-7} + \frac{8}{b-5}}{\frac{6}{b^2 - 12b + 35}}$$

$$37. \frac{3 - \frac{4}{a+4}}{\frac{5}{a+4} - \frac{6}{a-1}}$$

$$38. \frac{\frac{6}{x-5} + 7}{\frac{8}{x-5} - \frac{9}{x+3}}$$

$$39. \frac{\frac{y^2 - y - 6}{y^2 + 2y + 1}}{\frac{y^2 - 2y - 8}{y^2 + 7y + 6}}$$

$$40. \frac{\frac{a^2 + 3a - 10}{a^2 - 5a - 14}}{\frac{a^2 + 6a + 5}{a^2 - 8a + 7}}$$

Solve the following word problems.

41. A refrigeration coefficient-of-performance formula for the ideal refrigerator is given by

$$cp = \frac{1}{\frac{T_2}{T_1} - 1}$$

Simplify the right member.

42. In electronics, a formula for coupled inductance in parallel with fields aiding is given by

$$L_t = \frac{1}{\frac{1}{L_1 - M} + \frac{1}{L_2 + M}}$$

Simplify the right member.

43. In electronics, a formula for self-inductance of circuits in parallel is given by

$$L_t = \frac{1}{\frac{1}{L_1} + \frac{1}{L_2} + \frac{1}{L_3}}$$

Simplify the right member.

44. Coupled inductance with circuits connected in parallel with opposing fields is given by

$$L_t = \frac{1}{\frac{1}{L_1 - M} + \frac{1}{L_2 - M}}$$

Simplify the right member.

### Review exercises

Reduce the following expressions to lowest terms. See section 5-2.

1.  $\frac{36}{42}$

2.  $\frac{x^2 - 5x - 14}{x^2 + 4x + 4}$

3. Subtract  $(3x^3 - 2x^2 + x - 12) - (x^3 - 5x^2 + 9)$ . See section 2-3.

Solve the following equations. See sections 2-6 and 4-7.

4.  $4x + 3x = 21$

5.  $8y - 4 = 5y - 10$

6.  $x^2 + 2x - 3 = 0$

Determine the domain of the following rational expressions. See section 5-1.

7.  $\frac{3}{x + 7}$

8.  $\frac{y - 4}{y^2 - 4}$

9. Perform the indicated operations, if possible. See section 1-7.

a.  $\frac{4}{0}$

b.  $\frac{0}{-3}$

## 6-5 ■ Rational equations

### A rational equation

An algebraic equation that contains *at least one* rational expression is called a **rational equation**.

The basic operations for solving equations that you learned in chapter 2 will apply to rational equations once the denominators in the equation are eliminated. We accomplish this by using the multiplication property of equality. The multiplier is the least common denominator of all denominators in the rational expressions of the equation.

### Solving a rational equation

1. Find the LCD of all denominators.
2. Eliminate the denominators by multiplying each term of both members of the equation by the LCD of the denominators in the equation.
3. Use the four steps from section 2-6 to solve the resulting equation.

### Example 6-5 A

Find the solution set of each of the following rational equations.

$$1. \frac{x-3}{4} = \frac{x}{8}$$

We can see that the LCD of the denominators 4 and 8 is 8.

$$\frac{8}{1} \cdot \frac{(x-3)}{4} = \frac{8}{1} \cdot \frac{x}{8}$$

Multiply each member by 8

$$2(x-3) = x$$

Reduce in each member

$$2x - 6 = x$$

Multiply as indicated

$$x - 6 = 0$$

Subtract  $x$  from each member

$$x = 6$$

Add 6 to each member

The solution set is  $\{6\}$ . If we wish to check our work, we substitute 6 for  $x$  in the original equation to obtain the equivalent equation.

$$\frac{6-3}{4} = \frac{6}{8}$$

Replace  $x$  with 6 in the original equation

$$\frac{3}{4} = \frac{3}{4}$$

(True)

$$2. \frac{t}{4} - \frac{t-4}{5} = \frac{7}{10}$$

The LCD of the denominators 4, 5, and 10 is 20.

$$\frac{20}{1} \cdot \frac{t}{4} - \frac{20}{1} \cdot \frac{t-4}{5} = \frac{20}{1} \cdot \frac{7}{10}$$

Multiply each term by the LCD 20

$$5t - 4(t-4) = 2 \cdot 7$$

Reduce each term

$$5t - 4t + 16 = 14$$

Multiply as indicated

$$t + 16 = 14$$

Combine like terms

$$t = -2$$

Subtract 16 from each member

The solution set is  $\{-2\}$ . Check your answer by replacing  $t$  with  $-2$  in the original equation.

**Note** A common error that is made when multiplying  $-4(t-4)$  is to get  $-4t - 16$ . Do not forget that you are using the distributive property to multiply  $-4$  times each term in the group  $(t-4)$ . The correct result is  $-4t + 16$ .

$$3. \frac{5}{3a} + \frac{4}{9} = \frac{5}{12a}$$

We determine that the LCD of the denominators is  $36a$ .

$$36a \cdot \frac{5}{3a} + 36a \cdot \frac{4}{9} = 36a \cdot \frac{5}{12a} \quad (a \neq 0)$$

Multiply each term by the LCD  $36a$

$$12 \cdot 5 + 4a \cdot 4 = 3 \cdot 5$$

Reduce each term

$$60 + 16a = 15$$

Multiply as indicated

$$16a = -45$$

Subtract 60 from each member

$$a = -\frac{45}{16}$$

Divide each member by 16

The solution set is  $\left\{-\frac{45}{16}\right\}$ . Check your solution.



**Note** The domain of the variable is every real number except 0 since two of the terms are undefined when  $a = 0$ . This is an important observation to make as shown in example 4.

$$4. \frac{3}{y} = \frac{4}{y^2 - 2y} - \frac{2}{y - 2}$$

Factor the denominator  $y^2 - 2y$  to get  $y(y - 2)$ . We determine the LCD of the denominators to be  $y(y - 2)$ .

$$y(y - 2) \cdot \frac{3}{y} = y(y - 2) \cdot \frac{4}{y(y - 2)} - y(y - 2) \cdot \frac{2}{y - 2} \quad (y \neq 0, 2)$$

Multiply each term by the LCD  $y(y - 2)$

$$(y - 2) \cdot 3 = 4 - y \cdot 2$$

Reduce each term

$$3y - 6 = 4 - 2y$$

Multiply as indicated

$$3y + 2y = 4 + 6$$

Add 6 and  $2y$  to each member

$$5y = 10$$

Combine like terms

$$y = 2$$

Divide each member by 5

Two is *not* in the domain of the variable  $y$ , since  $y = 2$  makes the denominator  $y - 2 = 0$ . So 2 cannot be a solution of the equation. Therefore, the solution set is  $\emptyset$ .

We conclude there is no solution for the equation of example 4. The number 2 in that example is called an *extraneous solution*. An extraneous solution can occur whenever the variable appears in the denominator of one or more of the terms of the equation. Thus, you should *always* check your solution(s) of a rational equation when a variable appears in the denominator.

$$5. \frac{1}{3}x^2 - \frac{5}{2}x + 3 = 0$$

Multiply each term of the equation by 6, the LCD of the denominators 2 and 3.

$$6 \cdot \frac{1}{3}x^2 - 6 \cdot \frac{5}{2}x + 6 \cdot 3 = 6 \cdot 0$$

$$2x^2 - 15x + 18 = 0$$

Write the equation in standard form

$$(2x - 3)(x - 6) = 0$$

Factor  $2x^2 - 15x + 18 = (2x - 3)(x - 6)$

$$2x - 3 = 0 \quad \text{or} \quad x - 6 = 0$$

Set each factor equal to 0

$$2x = 3$$

$$x = 6$$

Solve each equation

$$x = \frac{3}{2}$$

$$x = 6$$

The solution set is  $\left\{\frac{3}{2}, 6\right\}$ . Check your solution.

**Note** We eliminate all the denominators *only* when solving rational equations. This is *never done* when adding or subtracting rational expressions.

To illustrate, given the equation  $\frac{6}{x} - \frac{4}{x^2} = 0$ , we multiply each term by  $x^2$ ,

whereas given the subtraction problem  $\frac{6}{x} - \frac{4}{x^2}$ , we do *not* multiply each term by  $x^2$ .

► **Quick check** Find the solution set of  $\frac{6}{4z} + \frac{7}{8} = \frac{9}{16z}$

### Rational equations in more than one variable

In scientific fields, equations and formulas involving rational expressions and *more than one variable* are common. It is often desirable to solve such equations for one variable in terms of the other variables in the equation. The procedures for finding such solutions are identical to those used in solving the preceding equations.

#### Solving a rational equation in more than one variable

1. Remove the fractions by multiplying each member of the equation by the LCD of the denominators in the equation.
2. Collect all terms containing the variable you are solving for in one member of the equation and all other terms in the other member.
3. Factor out the variable you are solving for if it appears in more than one term.
4. Divide each member by the coefficient of the variable for which you are solving.

### Example 6-5 B

Solve each rational equation for the indicated variable.

1. Solve  $\frac{a}{3} + \frac{3x}{2} = c$  for  $x$ .

Determine that the LCD of 3 and 2 is 6.

$$\begin{aligned}
 6 \cdot \frac{a}{3} + 6 \cdot \frac{3x}{2} &= 6 \cdot c && \text{Multiply each term by 6} \\
 2 \cdot a + 3 \cdot 3x &= 6c && \text{Reduce where possible} \\
 2a + 9x &= 6c && \text{Multiply in left member} \\
 2a + 9x - 2a &= 6c - 2a && \text{Subtract } 2a \text{ from each member} \\
 9x &= 6c - 2a && \text{Combine like terms} \\
 \frac{9x}{9} &= \frac{6c - 2a}{9} && \text{Divide each member by the coefficient 9} \\
 x &= \frac{6c - 2a}{9}
 \end{aligned}$$

2. Solve  $\frac{1}{a} = \frac{1}{b} + \frac{1}{c}$  for  $c$  ( $a, b, c \neq 0$ ).

Determine the LCD of  $a$ ,  $b$ , and  $c$  is  $abc$ .

$$\begin{aligned}
 abc \cdot \frac{1}{a} &= abc \cdot \frac{1}{b} + abc \cdot \frac{1}{c} && \text{Multiply each term by } abc \\
 bc &= ac + ab && \text{Reduce where possible}
 \end{aligned}$$

**Note** Since we are solving for  $c$ , we must get all terms containing  $c$  in the same member of the equation.

$$bc - ac = ac + ab - ac$$

Subtract  $ac$  from each member

$$bc - ac = ab$$

$$(b - a)c = ab$$

Factor  $c$  from each term in the left member

$$\frac{(b - a)c}{(b - a)} = \frac{ab}{(b - a)}$$

Divide each member by the coefficient of  $c$ ,  $(b - a)$ 

$$c = \frac{ab}{b - a}$$

Reduce in left member

► **Quick check** Solve  $\frac{5}{a} - \frac{3}{b} = 3$  for  $a$ .

### Mastery points

Can you

- Solve equations containing rational expressions?
- Solve rational equations for one variable in terms of the other variables?

### Exercise 6-5

Find the solution set of each rational equation. Indicate any restrictions on the domain of the variable. See example 6-5 A.

**Example**  $\frac{6}{4z} + \frac{7}{8} = \frac{9}{16z} \quad (z \neq 0)$

**Solution** The LCD of the denominators is  $16z$ .

$$16z \cdot \frac{6}{4z} + 16z \cdot \frac{7}{8} = 16z \cdot \frac{9}{16z}$$

Multiply each term by  $16z$ 

$$4 \cdot 6 + 2z \cdot 7 = 9$$

Reduce in each term

$$24 + 14z = 9$$

Multiply in each term

$$14z = -15$$

Subtract 24 from each member

$$z = -\frac{15}{14}$$

Divide each member by 14

The solution set is  $\left\{-\frac{15}{14}\right\}$ .

1.  $\frac{y}{4} = \frac{2}{3}$

2.  $\frac{4x}{5} - \frac{2}{3} = 4$

3.  $\frac{p}{6} = \frac{7}{9}$

4.  $\frac{a}{3} + \frac{5}{2} = 6$

5.  $\frac{z}{8} + 3 = \frac{1}{4}$

6.  $\frac{3R}{4} - 5 = \frac{5}{6}$

7.  $\frac{3a}{6} + \frac{2a}{5} = 1$

8.  $\frac{5x}{8} - \frac{x}{12} = 3$

9.  $\frac{2x + 1}{7} - \frac{2x - 3}{14} = 1$

10.  $\frac{3a + 1}{9} + \frac{1}{12} = \frac{2a - 1}{3}$

11.  $\frac{3}{2x} = \frac{4}{5} + \frac{2}{x}$

12.  $\frac{4}{b} - \frac{7}{3b} = \frac{2}{5}$



13.  $\frac{2}{3R} + \frac{3}{2R} + \frac{1}{R} = 4$

16.  $\frac{16}{5a} - 1 = 5 + \frac{3}{4a}$

19.  $\frac{3p+2}{7p} - 3 = \frac{p}{14p}$

22.  $\frac{R+2}{10R} + \frac{4R-1}{4R} = 2$

25.  $\frac{10}{3z+1} = \frac{3}{5}$

28.  $\frac{1}{x+5} = \frac{3}{x-5} - \frac{10}{x^2-25}$

31.  $\frac{5}{a^2-25} + \frac{3}{a-5} = \frac{4}{a+5}$

33.  $\frac{5}{x^2+x-6} = \frac{2}{x^2+3x-10}$

35.  $\frac{1}{a-3} + \frac{2}{a+4} = \frac{6}{a^2+a-12}$

37.  $3x^2 + 4x + \frac{4}{3} = 0$

38.  $3x^2 + \frac{11}{2}x + \frac{3}{2} = 0$

39.  $b^2 + \frac{3}{2}b = \frac{9}{2}$

40.  $\frac{2}{3}x^2 + x = \frac{20}{3}$

41.  $\frac{3}{4}z^2 = 2 - \frac{5}{2}z$

42.  $x^2 - \frac{5}{6}x = \frac{2}{3}$

43.  $x^2 = \frac{2}{3} - \frac{x}{3}$

44.  $x^2 + \frac{4}{3}x = -\frac{4}{9}$

14.  $\frac{3}{w} - \frac{6}{5w} + \frac{1}{2w} = 5$

17.  $\frac{5}{3b} - \frac{1}{2} = \frac{7}{6b}$

20.  $\frac{5-x}{8x} = \frac{2x+5}{6x}$

23.  $\frac{4}{x-4} = \frac{5}{x+4}$

26.  $\frac{3}{2a+3} = \frac{1}{7}$

29.  $\frac{b-1}{b^2-4} = \frac{6}{b+2}$

15.  $\frac{4}{6y} + 5 = \frac{1}{9y} + 2$

18.  $3 - \frac{5}{9x} = \frac{4}{6x}$

21.  $\frac{a-4}{3a} = \frac{2a-1}{4a}$

24.  $\frac{9}{3-x} = \frac{8}{2x+1}$

27.  $\frac{y}{y-2} - 2 = \frac{2}{y-2}$

30.  $\frac{5}{R+3} = \frac{4R+3}{R^2-9}$

32.  $\frac{6}{2p-4} + \frac{4}{p-2} = \frac{1}{p^2-4}$

34.  $\frac{8}{a^2-6a+8} = \frac{1}{a^2-16}$

36.  $\frac{5}{2x-1} - \frac{7}{3x+2} = \frac{9}{6x^2+x-2}$

Solve the following equations for the indicated letter. Assume all denominators are nonzero. See example 6-5 B.

**Example**  $\frac{5}{a} - \frac{3}{b} = 3$  for  $a$

**Solution**  $ab \cdot \frac{5}{a} - ab \cdot \frac{3}{b} = ab \cdot 3$

$$5b - 3a = 3ab$$

$$5b = 3ab + 3a$$

So  $5b = a(3b + 3)$

$$a = \frac{5b}{3b + 3}$$

Multiply each term by  $ab$

Reduce each term

Add  $3a$  to each member

Factor  $a$  in the right member

Divide each member by  $3b + 3$

45.  $\frac{2}{x} + \frac{1}{y} = 3$  for  $x$

46.  $\frac{5}{I} - 6 = \frac{8}{E}$  for  $I$

47.  $\frac{1}{c} = \frac{1}{c_1} + \frac{1}{c_2}$  for  $c_1$

48.  $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3}$  for  $R$

49.  $\frac{1}{8} = \frac{1}{a} + \frac{1}{b}$  for  $a$

50.  $\frac{1}{x} - \frac{1}{y} - \frac{1}{z} = 6$  for  $y$

51.  $\frac{3}{a} - \frac{4}{b} = \frac{5}{ab}$  for  $a$

52.  $\frac{5}{m} + 4 = \frac{6a}{2m} + 3b$  for  $m$

53.  $\frac{x+4}{2} + \frac{y-3}{5} = \frac{2}{10}$  for  $x$

Solve the following word problems.

54. The principal amount of money  $P$  in a savings account paying interest rate  $r$ , over a given period of time  $t$ , which pays interest  $I$  is given by  $P = \frac{I}{rt}$ . Solve the equation for  $r$ .
55. The pressure per square inch of steam or water in a pipe,  $p$ , is given by  $p = \frac{P}{LD}$ , where  
 $P$  = the total pressure on a diametral plane,  
 $L$  = the length of the pipe in inches, and  
 $D$  = the diameter of the pipe in inches.  
 Solve for  $D$ .
56. The safe internal unit pressure,  $p$ , in a given pipe of given thickness is given by  $p = \frac{2st}{D}$ , where  
 $s$  = the unit tensile stress,  
 $t$  = the thickness of the pipe, and  
 $D$  = the diameter of the pipe. Solve for  $s$ .
57. The pitch diameter of a gear,  $P$ , is given by  
 $P = \frac{D \cdot N}{N + 2}$ , where  $D$  = the outside diameter of the gear and  $N$  = the number of teeth in the gear.  
 Solve for  $N$ .
58. The rule governing the speeds of two gears, one the driver gear and the other the driven gear, is given by  $\frac{T_A}{T_B} = \frac{R_B}{R_A}$ , where  $T_A$  = the number of teeth in the driven gear,  $T_B$  = the number of teeth in the driver gear,  $R_B$  = the revolutions per minute of the driven gear, and  $R_A$  = the revolutions per minute of the driver gear. Solve for  $T_B$ .
59. Given  $F$  is the force on the large piston and  $f$  is the force on the small piston of a hydraulic press, then  $\frac{F}{f} = \frac{A}{a}$ , where  $A$  is the area of the large piston and  $a$  is the area of the small piston. Solve for  $f$ .
60. The kinetic energy of a body,  $KE$ , is computed by  $KE = \frac{Wv^2}{2g}$ , where  $v$  is the velocity expressed in feet per second,  $W$  is the weight in pounds, and  $g$  is acceleration due to gravity. Solve for  $W$ .
61. Charles' Law is in regard to the relationship between pressure  $P$ , volume  $V$ , and the absolute temperature  $T$  of a gas as it expands when heated. Charles' Law is represented by  $\frac{P_1V_1}{T_1} = \frac{P_2V_2}{T_2}$ , where  $P_1$ ,  $V_1$ , and  $T_1$  are the initial conditions of the gas and  $P_2$ ,  $V_2$ , and  $T_2$  are the final conditions. Solve for  $T_2$ .
62. The coefficient of linear expansion,  $k$ , of a solid when heated is given by  $k = \frac{L_t - L_0}{L_0 t}$ , where  $L_0$  is the length at  $0^\circ \text{C}$ ,  $L_t$  is the length at  $t^\circ \text{C}$ , and  $t$  is any given temperature in Celsius. Solve for  $L_t$ . Solve for  $L_0$ .
63. The coefficient of linear expansion  $\alpha$  (alpha) when metal is heated is given by  $\alpha = \frac{L_2 - L_1}{L_1(t_2 - t_1)}$ . Solve for  $L_1$ .
64. Carnot's ideal efficiency of any heat engine operating between the temperature limits  $T_1$  and  $T_2$  is given by  $CE = \frac{T_1 - T_2}{T_1}$ . Solve for  $T_1$ .
65. A formula for resistors in parallel is given by  $R = \frac{R_1R_2}{R_1 + R_2}$ . Solve for  $R_1$ .
66. In electronics, a formula for the unknown resistance  $R_x$  in a battery is given by  $R_x = R_m \left( \frac{E_1}{E_2} - 1 \right)$ . Solve for  $E_2$ .

## Review exercises

Evaluate each expression for the given values. See sections 2–2 and 5–1.

- $5x - 3y + z$  when  $x = 1$ ,  
 $y = -2$ , and  $z = 3$
- $\frac{3a - b}{2a + b}$  when  $a = 4$  and  
 $b = -5$
- $\frac{y_1 - y_2}{x_1 - x_2}$  when  $x_1 = 3$ ,  $x_2 = 1$ ,  
 $y_1 = 3$ , and  $y_2 = -5$



## Student Loans for up to **\$40,000** per year\*

Defer payments until after graduation.\*\*  
Fast preliminary approval, usually in minutes.

 **Apply Now!**

Go to [gmacbankfunding.com](http://gmacbankfunding.com)

## Apply online in as little as 15 minutes

- Loans up to \$40,000 per academic year\*
- Good for tuition and other educational expenses: books, fees, a laptop, room and board, travel home, etc.
- Get a check in as few as 5 business days
- Start payments now or up to six months after graduation\*\*
- Reduce your interest rate by as much as 0.50% with automatic payments\*\*\*

All loans are subject to application and credit approval.

\* Undergraduate and graduate borrowers may borrow annually up to the lesser of the cost of attendance or \$30,000 (\$40,000 for certain schools where it has been determined that the annual cost of attendance exceeds \$30,000). Borrowers in the Continuing Education loan program may borrow annually up to \$30,000.

\*\* Undergraduate students may choose to defer repayment until six months after graduation or ceasing to be enrolled at least half time in school. Interest only and immediate repayment options also available.

\*\*\* A 0.25% interest rate reduction is available for borrowers who elect to have monthly principal and interest payments transferred electronically from a savings or checking account. The interest rate reduction will begin when automatic principal and interest payments start, and will remain in effect as long as automatic payments continue without interruption. This reduced interest rate will return to contract rate if automatic payments are cancelled, rejected or returned for any reason. Upon request, borrowers are also entitled to an additional 0.25% interest rate reduction if (1) the first 36 payments of principal and interest are paid on time, and (2) at any time prior to the 36th on time payment, the borrower who receives the monthly bill elects to have monthly principal and interest payments transferred electronically from a savings or checking account, and continues to make such automatic payments through the 36th payment. This reduced interest rate will not be returned to contract rate if, after receiving the benefit, the borrower discontinues automatic electronic payment. The lender and servicer reserve the right to modify or discontinue borrower benefit programs (other than the co-signer release benefit) at any time without notice.

GMAC Bank Member FDIC



Solve each equation for  $y$ . See section 2-7.

4.  $5x + y = 4$

5.  $2x + 3y = 6$

6.  $x - 4y = 8$

Simplify the following expressions. Assume all denominators are nonzero. Answer with positive exponents only. See section 3-4.

7.  $(3y^{-1})(y^2x^{-2})$

8.  $\frac{x^{-3}y^3}{x^2y^{-1}}$

## 6-6 ■ Rational expression applications

Rational expressions occurring in rational equations have many applications in the physical and scientific world. We now wish to discuss some of the more common applications.

### ■ Example 6-6 A

Choose a variable, set up an appropriate equation, and solve the following problems.

1. A water holding tank is fed by two pipes. If it takes the smaller pipe 12 hours to fill the tank and the larger pipe 9 hours to fill the tank, how long would it take to fill the tank if both pipes are open? (This is called a *work* problem.)

Let  $x$  = the number of hours it takes the two pipes to fill the tank. Now,

- a. the smaller pipe can fill  $\frac{1}{12}$  of the tank in 1 hour,
- b. the larger pipe can fill  $\frac{1}{9}$  of the tank in 1 hour,
- c. the two pipes together can fill  $\frac{1}{x}$  of the tank in 1 hour.

The amount of the tank filled by the smaller pipe in 1 hour plus the amount of the tank filled by the larger pipe in 1 hour must be equal to the amount of the tank filled by the two pipes together in 1 hour. Thus, the equation is

$$\begin{array}{c} \text{Amount by larger pipe} \\ \downarrow \\ \text{Amount by smaller pipe} \rightarrow \frac{1}{12} + \frac{1}{9} = \frac{1}{x} \leftarrow \text{Amount together} \end{array}$$

Multiply each term by the LCM of 12, 9, and  $x$ , which is  $36x$ .

$$\begin{array}{rcl} 36x \cdot \frac{1}{12} + 36x \cdot \frac{1}{9} & = & 36x \cdot \frac{1}{x} \\ 3x + 4x & = & 36 \\ 7x & = & 36 \\ x & = & \frac{36}{7} \end{array} \quad \begin{array}{l} \text{Reduce in each term} \\ \text{Combine in left member} \\ \text{Divide each member by 7} \end{array}$$

Therefore, together the two pipes can fill the tank in  $\frac{36}{7}$  hours, or  $5\frac{1}{7}$  hours.

**Note** That is, approximately 5 hours and 9 minutes.

2. Jim James drove a distance of 120 miles, part at 50 miles per hour (mph) and part at 60 mph. If he drove the 120 miles in  $2\frac{1}{4}$  hours, how many miles did he drive at 50 mph? (This is called a *time–rate–distance* problem.)

**Note** In a distance ( $d$ )–rate ( $r$ )–time ( $t$ ) problem, we use  $d = rt$ ,  $t = \frac{d}{r}$ , or  $r = \frac{d}{t}$ .

Let  $x$  = the distance he drove at 50 mph. Then  $120 - x$  = the distance he drove at 60 mph.

The time traveled at 50 mph plus the time traveled at 60 mph equals the total time traveled,  $2\frac{1}{4}$  hr (or  $\frac{9}{4}$  hr). We use the following table for time–rate–distance problems.

distance ( $d$ )	rate ( $r$ )	time ( $t$ )
$x$	50	$\frac{x}{50}$
$120 - x$	60	$\frac{120 - x}{60}$

$t = \frac{d}{r}$

The equation is then

$$\begin{array}{l}
 \text{Time at 60 mph} \\
 \downarrow \\
 \text{Time at 50 mph} \rightarrow \frac{x}{50} + \frac{120 - x}{60} = \frac{9}{4} \leftarrow \text{Total time} \\
 300 \cdot \frac{x}{50} + 300 \cdot \frac{120 - x}{60} = 300 \cdot \frac{9}{4} \quad \text{Multiply each term by the LCD 300} \\
 6x + 5(120 - x) = 675 \quad \text{Reduce in each term} \\
 6x + 600 - 5x = 675 \quad \text{Distributive property} \\
 x + 600 = 675 \quad \text{Combine in left member} \\
 x = 75 \quad \text{Subtract 600 from each member}
 \end{array}$$

Thus, Jim James drove 75 miles at 50 mph.

**Note** He drove  $120 - x = 120 - 75 = 45$  miles at 60 mph.

3. The denominator of a fraction is 3 more than the numerator. If 4 is added to the numerator and the denominator, the resulting fraction is  $\frac{3}{4}$ . Find the original fraction.

Let  $x$  = the numerator of the original fraction. Then  $x + 3$  = the denominator of the original fraction. The equation we get is

$$\frac{x + 4}{(x + 3) + 4} = \frac{3}{4}$$

then

$$\frac{x+4}{x+7} = \frac{3}{4}$$

The LCD of 4 and  $x+7$  is  $4(x+7)$ .

$$4(x+7) \cdot \frac{x+4}{x+7} = 4(x+7) \cdot \frac{3}{4} \quad (x \neq -7) \quad \text{Multiply each term by } 4(x+7)$$

$$4(x+4) = (x+7) \cdot 3 \quad \text{Reduce each term}$$

$$4x + 16 = 3x + 21 \quad \text{Multiply as indicated}$$


$$x = 5 \quad \text{Subtract } 3x \text{ and } 16 \text{ from each member}$$

$$\text{and } x+3 = 8$$

The original fraction is  $\frac{5}{8}$ .

4. In an electrical circuit, when two resistors are connected in parallel, the total resistance  $R$  of the circuit in ohms is given by

$$\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$$



where  $R_1$  and  $R_2$  are the resistances of the two resistors in ohms and the circuit is connected in parallel as shown in the diagram. Find the total resistance  $R$  of an electrical circuit having two resistors connected in parallel if their resistances are 4 ohms and 6 ohms.

We want  $R$  if  $R_1 = 4$  ohms and  $R_2 = 6$  ohms.

$$\frac{1}{R} = \frac{1}{4} + \frac{1}{6}$$

Multiply both members by the LCM of 4, 6, and  $R$ , which is  $12R$ .

$$12R \cdot \frac{1}{R} = 12R \cdot \frac{1}{4} + 12R \cdot \frac{1}{6} \quad (R \neq 0)$$

$$12 = 3R + 2R$$

$$12 = 5R$$

$$\frac{12}{5} = R$$

Therefore the total resistance is  $\frac{12}{5}$ , or  $2\frac{2}{5}$ , ohms.

► **Quick check** If the same number is added to the numerator and the denominator of  $\frac{3}{4}$ , the result is the fraction  $\frac{5}{6}$ . What is the number?

### Mastery points

Can you

- Solve work problems?
- Solve distance-rate-time problems?
- Solve number problems?
- Solve resistance in electric circuits problems?



## Exercise 6-6

Choose a variable, set up an equation, and solve the following problems.

**Example** If the same number is added to the numerator and the denominator of  $\frac{3}{4}$ , the result is the fraction  $\frac{5}{6}$ . What is the number?

**Solution** Let  $x$  = the number to be added to the numerator and the denominator. The equation is then

$$\begin{aligned}\frac{3+x}{4+x} &= \frac{5}{6} \\ 6(4+x) \frac{3+x}{4+x} &= 6(4+x) \cdot \frac{5}{6} && \text{Multiply each member by the LCD } 6(4+x) \\ 6(3+x) &= (4+x) \cdot 5 && \text{Reduce in each member} \\ 18+6x &= 20+5x && \text{Multiply as indicated} \\ x &= 2 && \text{Solve for } x\end{aligned}$$

The number to be added is 2. (Check:  $\frac{3+2}{4+2} = \frac{5}{6}$ )

See example 6-6 A-1.

1. Jim can mow his parents' lawn in 50 minutes. His younger brother Kenny can mow the lawn in 70 minutes. How long would it take for the boys to mow the lawn if they mow together?
2. In a factory, worker  $A$  can do a certain job in 6 hours while worker  $B$  can do the same job in 5 hours. How long would it take workers  $A$  and  $B$  to do the same job working together?
3. Three different sized pipes feed water into a swimming pool. If the pipes can fill the same pool individually in 6 hours, 8 hours, and 9 hours, respectively, how long would it take to fill the same pool if all three pipes were open?
4. During "clean-up week" in Podunk Junction, Jane, Joan, and Ruth work to clean up a vacant lot. If the girls can do the job individually in 2 hours, 3 hours, and 4 hours, respectively, how long will it take them, working together, to clean the lot?
5. It takes two men—Harry and Dick—working together 4 hours to paint the exterior of a house. If Harry could do the job in 6 hours working alone, how long would it take Dick to paint the house alone?
6. A water tank has two drain pipes. If the larger pipe could empty the tank in 45 minutes and the two pipes together could drain the tank in 30 minutes, how long would it take for the smaller pipe to drain the tank?
7. Three combines— $A$ ,  $B$ , and  $C$ —working together can harvest a field of oats in  $1\frac{1}{2}$  hours. If  $A$  could do the job alone in 5 hours and  $B$  could do it alone in 6 hours, how long would it take for combine  $C$  to do the same job alone?
8. In a pizzeria, three women are at work making pizzas. The three could make 50 pizzas in  $1\frac{1}{2}$  hours working together. If two of the women could make all the pizzas in 5 hours and 6 hours, respectively, working alone, how long would it take the third woman to make the 50 pizzas working alone?
9. A tank has one inlet pipe and one outlet pipe. If the inlet pipe can fill the tank in  $3\frac{1}{3}$  hours and the outlet pipe can empty the tank in 5 hours, how long would it take to fill the tank if both pipes are left open? (*Hint*: Subtract the drainage.)
10. A sink drain, when left open, can empty a sink full of water in 4 minutes. If the cold water and hot water faucets can, when fully open, fill the sink in  $2\frac{1}{2}$  minutes and 3 minutes, respectively, how long would it take to fill the sink if all three are open simultaneously?



See example 6-6 A-2.

11. Jack drove 320 miles in  $5\frac{1}{2}$  hours. If he drove part of the trip averaging 55 mph and the rest averaging 60 mph, how many miles did he drive at each speed?
12. A. J. Foyt, when driving the Indianapolis 500 Race, averaged 198 mph over part of the race. Due to an accident on the track, he averaged 160 mph over the rest. If the race took 2 hours and 40 minutes to run, how many miles did he drive at an average of 198 mph? (Round the answer to the nearest integer.)
13. Car *A* travels 120 miles in the same time that car *B* travels 150 miles. If car *B* averages 10 mph faster than car *A*, how fast is each car traveling?
14. A freight train travels 260 kilometers in the same time a passenger train travels 320 kilometers. If the passenger train averages 15 kilometers per hour faster than the freight train, what was the average speed of the freight train?
15. Sheila can row a boat 2 mph in still water. How fast is the current of a river if she takes the same length of time to row 4 miles upstream as she does to row 10 miles downstream? (*Hint*: Subtract current upstream and add downstream.)
16. An airplane flew 1,000 miles with the wind in the same length of time it took to fly 850 miles against the wind. If the wind was blowing at 25 miles per hour, what was the average air speed of the plane?
17. On a trip from Detroit to Columbus, Ohio, Mrs. Smith drove at an average speed of 60 mph. Returning, her average speed was 55 mph. If it took her  $\frac{1}{3}$  hour longer on the return trip, how far is it from Detroit to Columbus?
18. A jet plane flew at an average speed of 240 miles per hour going from city *A* to city *B* and averaged 300 miles per hour on the return flight. Its return flight took 1 hour and 40 minutes less time. How far is it from city *A* to city *B*? (Disregard any wind.)

See example 6-6 A-3.

19. The numerator of a given fraction is 4 less than the denominator. If 5 is added to both the numerator and the denominator, the resulting fraction is  $\frac{5}{7}$ .  
What is the original fraction?
20. The denominator of a fraction exceeds the numerator by 7. If 3 is added to the numerator and 1 is subtracted from the denominator, the resulting fraction is  $\frac{4}{5}$ . Find the original fraction.
21. One number is four times another number. The sum of their reciprocals is  $\frac{5}{12}$ . What are the numbers?
22. One number is four times another number. The sum of their reciprocals is  $\frac{1}{4}$ . What are the numbers?
23. If  $\frac{1}{2}$  is added to three times the reciprocal of a number, the result is 1. Find the number.
24. If  $\frac{1}{2}$  is subtracted from four times the reciprocal of a number, the result is 0. Find the number.
25. When a certain number is added to the numerator and subtracted from the denominator of the fraction  $\frac{2}{5}$ , the result is 6. What is the number?
26. When a certain number is added to the numerator and subtracted from the denominator of the fraction  $\frac{5}{9}$ , the result is  $\frac{4}{7}$ . What is the number?
27. A prescription for an illness calls for a child's dosage to be  $\frac{3}{5}$  of the number of pills that an adult takes. Together they use 24 pills. How many pills are taken by each?
28. An apprentice electrician receives  $\frac{3}{8}$  of the hourly wage of a journeyman electrician. Together their hourly wage is \$29.70. Find the hourly wage of each.
29. Pat and Mike earn a total of \$65.50 per week delivering papers. If Pat's earnings are  $\frac{5}{7}$  of Mike's, how much does Mike earn each day?



Use the formula  $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2}$  to solve the following exercises. See example 6-6 A-5.

- 30.** Two resistors in an electric circuit have resistances of 6 ohms and 8 ohms and are connected in parallel. Find the total resistance of the circuit.
- 31.** Two resistors of an electric circuit are connected in parallel. If one has a resistance of 5 ohms and the other has a resistance of 12 ohms, what is the total resistance in the circuit?
- 32.** Three resistors connected in parallel have resistances of 4 ohms, 6 ohms, and 10 ohms. What is the total resistance in the electric circuit?  
 (Hint: Use  $\frac{1}{R} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3}$ .)
- 33.** The total resistance in a parallel wiring circuit is 12 ohms. If the resistance in one branch is 30 ohms, what is the resistance in the other branch?
- 34.** The resistance in one branch of a two-resistor parallel wiring circuit is 10 ohms. If the total resistance in the circuit is 6 ohms, what is the resistance in the other branch?
- 35.** A three-resistor parallel wiring circuit has a total resistance of 10 ohms. If two of the branches of the circuit have resistances of 20 ohms and 30 ohms, what is the resistance in the third branch? (See hint in exercise 32.)

### Review exercises

Find the solution set of the following equations. See sections 2-6 and 2-7.

1.  $2y + 3 = 4y - 1$

2.  $2y - 3x = 6$  for  $y$

Factor the following expressions. See sections 4-2, 4-3, and 4-4.

3.  $8y^2 - 32$

4.  $x^2 + 20x + 100$

5.  $3y^2 - 4y - 4$

Combine the following rational expressions. See section 6-3.

6.  $\frac{3x}{x-1} + \frac{2x}{x+3}$

7.  $\frac{4y}{2y+1} - \frac{3y}{y-5}$

### Chapter 6 lead-in problem

Marc owns  $\frac{5}{8}$  interest in a print shop and his uncle owns  $\frac{1}{4}$  interest in the shop. In a given year, they shared earnings of \$140,000. How much did the shop earn that year?

#### Solution

Let  $x$  = the total earnings of the print shop.

	Interest	Shop earnings	Income
Marc	$\frac{5}{8}$	$x$	$\frac{5}{8}x$
Uncle	$\frac{1}{4}$	$x$	$\frac{1}{4}x$

Then  $\frac{5}{8}x + \frac{1}{4}x = 140,000$

$8 \cdot \frac{5}{8}x + 8 \cdot \frac{1}{4}x = 8 \cdot 140,000$

$5x + 2x = 1,120,000$

$7x = 1,120,000$

$x = 160,000$

Together they earned \$140,000

Multiply each term by the LCM of 4 and 8, which is 8

Perform indicated multiplications

Combine like terms

Divide each member by 7



## Chapter 6 summary

1. To *multiply* two rational expressions, we multiply the numerators and place that product over the product of the denominators.
2. To *divide* two rational expressions,  $\frac{P}{Q} \div \frac{R}{S}$ , we multiply  $\frac{P}{Q}$  by the reciprocal of  $\frac{R}{S}$ , which is  $\frac{S}{R}$ .
3. To find the *least common denominator* (LCD) of two or more rational expressions, we
  - a. write each denominator in completely factored form;
  - b. take each different factor that appears in the factorizations in step a and write them as a product;
  - c. raise each factor of step b to the greatest power it has in step a.
4. To obtain an *equivalent rational expression*, we multiply, or divide, the numerator and the denominator of the given rational expression by a *common factor*.
5. To *add* or *subtract* rational expressions, we must
  - a. write each rational expression as an equivalent rational expression with the same denominator, preferably the LCD;
  - b. add, or subtract, the numerators and place this sum, or difference, over the common denominator.
6. A **complex fraction** is a fraction whose numerator or denominator, or both, contains at least one fraction.

7. In the *complex fraction*

$$\frac{\frac{a}{b}}{\frac{c}{d}}$$

we call  $\frac{a}{b}$  the *primary numerator*,  $\frac{c}{d}$  the *primary denominator*, and  $b$  and  $d$  the *secondary denominators*.

8. To *simplify a complex fraction*, we
  - a. divide the primary numerator by the primary denominator after making a single fraction of each; or
  - b. multiply the primary numerator and the primary denominator by the LCM of the secondary denominators and reduce the result, where possible.
9. A **rational equation** is an equation that contains at least one rational expression.
10. To *solve a rational equation*, we
  - a. multiply each term of the equation by the LCD to eliminate the fractions;
  - b. use the basic procedures for solving equations;
  - c. check for extraneous solutions.

## Chapter 6 error analysis

1. Subtracting rational expressions

Example:  $\frac{3x-1}{x-2} - \frac{x+2}{x-2} = \frac{3x-1-x+2}{x-2}$   
 $= \frac{2x+1}{x-2}$

Correct answer:  $\frac{2x-3}{x-2}$

What error was made? (see page 240)

2. Solving rational equations

Example: Find the solution set of  $\frac{y}{3} - \frac{y+1}{4} = \frac{2}{12}$

$$12 \cdot \frac{y}{3} - 12 \cdot \frac{y+1}{4} = 12 \cdot \frac{2}{12}$$

$$4y - 3(y+1) = 2$$

$$4y - 3y + 3 = 2$$

$$y = -1 \quad \{-1\}$$

Correct answer:  $\{5\}$

What error was made? (see page 259)

3. Adding and subtracting rational expressions

Example:  $\frac{3}{x-5} + \frac{5}{5-x} = \frac{3+5}{x-5} = \frac{8}{x-5}$

Correct answer:  $\frac{-2}{x-5}$

What error was made? (see page 241)

4. Adding and subtracting rational expressions

Example:  $\frac{3}{x^2} + \frac{2}{x^2} = x^2 \cdot \frac{3}{x^2} + x^2 \cdot \frac{2}{x^2} = 3 + 2 = 5$

Correct answer:  $\frac{5}{x^2}$

What error was made? (see page 239)

5. Exponents

Example:  $(-4)^2 = -16$

Correct answer: 16

What error was made? (see page 57)

6. Order of operations

Example:  $3^2 + 12 \cdot 2 - 8 \div 2 = 17$

Correct answer: 29

What error was made? (see page 57)

7. Combining polynomials

Example:  $(4a^2b - 2ab^2) - (a^2b + ab^2) = 3a^2b - ab^2$

Correct answer:  $3a^2b - 3ab^2$

What error was made? (see page 82)

8. Solving linear inequalities

Example: If  $4 < -2x \leq 6$ , then  $-2 < x \leq -3$ .

Correct answer: If  $4 < -2x \leq 6$ , then  $-2 > x \geq -3$ .

What error was made? (see page 116)

## 9. Multiplication of like bases

Example:  $5 \cdot 5^3 = 5^3$ Correct answer:  $5^4$ 

What error was made? (see page 129)

## 10. Squaring a binomial

Example:  $(x - 6)^2 = (x)^2 - (6)^2 = x^2 - 36$ Correct answer:  $x^2 - 12x + 36$ 

What error was made? (see page 135)

**Chapter 6 critical thinking**

Which integers can be written as the sum of three consecutive integers?

Example:  $6 = 1 + 2 + 3$ **Chapter 6 review****[6-1]**

Perform the indicated multiplication and reduce the product to lowest terms.

1.  $\frac{15}{8a} \cdot \frac{12a}{5}$

2.  $\frac{24b}{7a} \cdot \frac{21a^2}{8b^2}$

3.  $\frac{x - 3y}{2x + y} \cdot \frac{2x - y}{x - 3y}$

4.  $\frac{5x - 10}{x + 3} \cdot \frac{3x + 9}{15}$

5.  $\frac{m^2 - n^2}{14x} \cdot \frac{35x^2}{5m + 5n}$

6.  $\frac{y}{y^2 - 1} \cdot \frac{y + 1}{y^2 - y}$

7.  $\frac{b^2 - 36}{b^2 + 12b + 36} \cdot \frac{2b + 12}{b - 6}$

8.  $\frac{5 - x}{6a - 3} \cdot \frac{2a - 1}{x^2 - 25}$

9.  $\frac{x^2 + x - 2}{x^2 - 4x - 12} \cdot \frac{x^2 - 7x + 6}{x^2 - 1}$

Find the indicated quotients and state the answer reduced to lowest terms. Assume all denominators are nonzero.

10.  $\frac{14a}{9} \div \frac{7}{3}$

11.  $\frac{10}{9b} \div \frac{35}{12b^2}$

12.  $\frac{24ab}{7} \div \frac{16a^2b^2}{21}$

13.  $\frac{2x - 1}{3x + 4} \div \frac{3x}{7x}$

14.  $\frac{x + 6}{x^2 - 4} \div \frac{(x + 6)^2}{x + 2}$

15.  $\frac{4a - 8}{2a + 1} \div \frac{6a - 12}{10a + 5}$

16.  $\frac{x^2 + 16x + 64}{x^2 + 9x + 8} \div \frac{x^2 - 64}{x + 1}$

17.  $\frac{1 - b}{(b + 3)^2} \div \frac{b^2 - 2b + 1}{b^2 - 9}$

18.  $\frac{9a^2 + 15a + 6}{a^2 + 3a - 4} \div \frac{36a^2 - 16}{3a^2 + 10a - 8}$

**[6-2]**

Find the least common denominator of rational expressions having the given denominators. Assume all denominators are nonzero.

19.  $9x^2$  and  $12x$

20.  $14ab$  and  $21a^2b^2$

21.  $x^2 - 9$  and  $2x + 6$

22.  $y^2 - 2y - 15$  and  $y^2 - 25$

23.  $4z - 8$ ,  $z^2 - 4$ , and  $z + 1$

24.  $x^2 + x$ ,  $x^2 + 2x + 1$ , and  $3x^2 - 2x - 5$

Find the indicated sum or difference. Assume all denominators are nonzero.

25.  $\frac{4}{x} + \frac{1}{-x}$

26.  $\frac{y}{y - 2} - \frac{3y}{2 - y}$

27.  $\frac{3x - 2}{3x - 1} - \frac{x + 6}{1 - 3x}$

28.  $\frac{x + 7}{2x - 5} + \frac{2x - 6}{5 - 2x}$

**[6-3]**

Add or subtract as indicated and reduce the answer to lowest terms. Assume all denominators are nonzero.

29.  $\frac{4x}{15} + \frac{8x}{21}$

30.  $\frac{25}{16a} - \frac{13}{12a}$

31.  $\frac{5}{3x + 1} + \frac{9}{4x - 3}$

32.  $\frac{4x}{x + 4} - \frac{7x}{x^2 - 16}$

33.  $\frac{4}{ab^2} + \frac{12}{5a^2b} - \frac{3}{4ab}$

34.  $4a + \frac{6}{a + 1}$



35.  $\frac{7}{x^2 + 1} - 10$

36.  $\frac{x+1}{4x} - \frac{3x-5}{8x^2} + \frac{5x+4}{12x}$

37.  $\frac{4y}{y^2 - 7y - 18} + \frac{9y}{y^2 - 4}$

38.  $\frac{5a}{x+3} + \frac{2}{x^2-9} - \frac{7a}{x-3}$

39.  $\frac{9}{x} - \frac{4}{x-5} + \frac{3}{x+4}$

40.  $\frac{5}{x^2 + 2xy + y^2} + \frac{5}{x^2 - xy - 2y^2}$

**[6-4]**

Simplify the given complex fractions and reduce to lowest terms.

41.  $\frac{\frac{4}{7}}{\frac{9}{4}}$

42.  $\frac{\frac{3}{4} - \frac{1}{2}}{\frac{1}{4} + \frac{3}{2}}$

43.  $\frac{\frac{4}{5} + 1}{2 - \frac{1}{5}}$

44.  $\frac{\frac{1}{x} - \frac{1}{y}}{\frac{1}{x} + \frac{1}{y}}$

45.  $\frac{\frac{4}{x^2} + \frac{3}{x}}{\frac{2}{x^2} - \frac{5}{x}}$

46.  $\frac{\frac{a}{b} - \frac{b}{a}}{\frac{a}{b} + \frac{b}{a}}$

47.  $\frac{\frac{1}{x} - \frac{1}{y}}{\frac{1}{xy}}$

48.  $\frac{\frac{1}{x} - \frac{1}{y}}{\frac{1}{x^2} - \frac{1}{y^2}}$

49.  $\frac{\frac{a^2}{b^2} - 2a - 3}{\frac{a - 3b}{ab}}$

**[6-5]**

Find the solution set of the following rational equations.

50.  $\frac{x}{8} - 3 = \frac{2x}{12} + 1$

51.  $\frac{4a+1}{4} + \frac{5}{6} = 3a - \frac{5}{8}$

52.  $\frac{12}{4a} - \frac{5}{6a} = 4$

53.  $\frac{7}{a+1} - \frac{1}{a^2-1} = \frac{2}{a+1}$

54.  $\frac{1}{y+3} - \frac{6}{y} = \frac{7}{y} + \frac{2}{y+3}$

55.  $\frac{7}{2c^2} + \frac{5}{6c^2} = \frac{7}{12c^3}$

56.  $\frac{x^2}{4} - 3x = 0$

57.  $\frac{2}{3}y^2 = \frac{3}{2}$

58.  $\frac{x^2}{2} - \frac{15}{2} = -x$

59.  $\frac{a^2}{3} = \frac{5a}{2} - 3$

Solve for the indicated variable.

60.  $\frac{a}{x} + \frac{b}{x} = 3$  for  $x$

61.  $\frac{y}{x+1} = \frac{y^2}{x-3}$  for  $x$

62.  $\frac{3}{4-y} = \frac{a}{b}$  for  $y$

63.  $\frac{4}{y-c} - \frac{5}{y+b} = 0$  for  $y$

Solve the following word problems.

64. An equation for tensile and compressional stresses is given by
- $y = \frac{Fl}{Ae}$
- . Solve for
- $l$
- .

65. The efficiency of a screw jack is calculated by the formula
- $E = \frac{Wp}{2 + LF}$
- . Solve for
- $L$
- .

66. A formula for the theoretical mechanical advantage,
- $M$
- , of a chain fall is given by

$$M = \frac{2R}{R-r}$$
 Solve for  $R$ .

67. The total reaction force
- $F$
- of air against a plane at the bottom of a vertical loop in centripetal force is given by

$$F = \frac{mv^2}{r} + m \cdot g$$
 Solve for  $m$ .

**[6-6]**

68. With different equipment, one painter can paint a house three times faster than a second painter. Working together they can do it in 4 hours. How long would it take each of them to paint the house working alone?

69. Paul can row his boat at a rate of 4 miles per hour in still water. It takes him as long to row 20 miles downstream as it takes him to row 8 miles upstream. What is the rate of the current?

70. The sum of three times a number and twice its reciprocal is 5. Find the number.



**Chapter 6 cumulative test**

Compute and simplify.

[1-5] 1.  $-\frac{1}{3} + \frac{1}{4} + (-6) + \frac{1}{6}$

[1-6] 2.  $-\frac{4}{5} \left( -\frac{15}{16} \right)$

[1-5] 3.  $\frac{5}{6} - \left( -\frac{3}{5} \right)$

[1-7] 4.  $-\frac{24}{35} \div \left( -\frac{3}{7} \right)$

Simplify the following expressions with only positive exponents.

[3-3] 5.  $x^8 \cdot x^3 \cdot x^0$

[3-3] 6.  $\frac{y^{-5}}{y^4}$

[3-4] 7.  $(-5x^2y^{-3})^2$

[2-3] 8.  $(9x^3 - 3x^2 + 4x - 8) - (-x^3 + x^2 - 7x + 1)$

Find the solution set of the following equations and inequalities.

[2-6] 9.  $-5(2x + 5) = -3x + 1$

[2-6] 10.  $\frac{1}{3}x - \frac{3}{4} = 6$

[4-7] 11.  $2y^2 - y = 6$

[2-9] 12.  $3x - 4 \leq 5x + 6$

Completely factor each expression.

[4-4] 13.  $2x^2 - 18$

[4-1] 14.  $6x^5 - 36x^3 + 9x^2$

[4-3] 15.  $4x^2 + 16x + 15$

[4-4] 16.  $3 - 12x^6$

Perform the indicated operations.

[3-2] 17.  $(7x - 6)^2$

[3-2] 18.  $\left( 5 - \frac{1}{2}x \right) \left( 5 + \frac{1}{2}x \right)$

[3-2] 19.  $(2x - 3)(3x^2 - 5x + 11)$

[3-2] 20.  $(3 - 2x^2)(5 - 4x^2)$

[5-4] 21. Find  $x$  when  $\frac{x}{24} = \frac{5}{7}$ .

[5-4] 22. Power (in foot-pounds per minute) is given by the ratio of work,  $w$ , to time,  $t$ . Find the power exerted by pushing an 80-pound load 180 feet in 6 minutes. (Hint:  $w = \text{weight} \cdot \text{length}$ .)

Perform the indicated operations and reduce the answers to lowest terms.

[6-1] 23.  $\frac{4a}{5b} \cdot \frac{a}{b}$

[6-1] 24.  $\frac{y+3}{y-6} \cdot \frac{y+7}{y-2}$

[6-1] 25.  $\frac{y^2-4}{y+3} \cdot \frac{y^2-9}{y+2}$

[6-1] 26.  $\frac{x^2-y^2}{9} \div \frac{y^2+xy}{9x-9}$

[6-3] 27.  $\frac{2}{x^2y} + \frac{4}{xy^2}$

[6-3] 28.  $\frac{6}{x-y} - \frac{3}{x+y}$

[6-3] 29.  $\frac{a+1}{a-2} + \frac{a-8}{2-a}$

[6-3] 30.  $\frac{7}{x^2-49} + \frac{6}{x^2-5x-14}$

[6-3] 31.  $\frac{2x}{x^2+x-6} - \frac{3x}{x^2-3x-18}$

Simplify the following complex rational expressions.

[6-4] 32.  $\frac{3 - \frac{1}{y}}{4 + \frac{5}{y}}$

[6-4] 33.  $\frac{\frac{1}{x} + \frac{1}{y}}{\frac{3}{y} - \frac{4}{x}}$

[6-4] 34.  $\frac{\frac{x-4}{x+7}}{\frac{x+5}{x^2-49}}$

Find the solution set of the following equations.

[6-5] 35.  $\frac{4}{x} = \frac{6}{2x-1}$

[6-5] 36.  $x^2 - 2 = \frac{17x}{3}$

[6-5] 37. Solve the equation  $\frac{1}{f} = \frac{1}{p} + \frac{1}{q}$  for  $q$ .

Choose a variable, set an appropriate equation, and solve the following problems.

- [6-6] 38. John can build a fence in 6 days and Harry can build the same fence in 4 days. How long would it take them to build the fence working together?

- [6-6] 39. The denominator of a fraction is two times the numerator. If 2 is added to the numerator and 2 is subtracted from the denominator, the resulting fraction is  $\frac{5}{7}$ . Find the original fraction.

15.  $\frac{R-4}{3R-1}$  16.  $\frac{5-n}{2+n}$  17.  $-8x^2$  18.  $2a-3+5a^2$   
 19.  $5x-3y^3+xy$  20.  $2a+3b-6ab^6$   
 21.  $4a+1+\frac{-2}{2a-1}$  22.  $3a-2+\frac{1}{a-5}$  23.  $x-7$   
 24.  $5x^2-x-4$  25.  $\frac{3}{7}$  or 3:7 26.  $\frac{9}{4}$  or 9:4 27.  $\frac{2}{5}$  or 2:5  
 28. 30  $\frac{\text{miles}}{\text{gallon}}$  or 30 miles per gallon 29.  $\frac{2,193}{881}$  or 2,193:881  
 30.  $\frac{7}{3}$  or 7:3 31.  $x=32$  32.  $a=3.6$  33.  $y=\frac{54}{5}$   
 34.  $p=\frac{2}{5}$  35. 35 feet 36. 12 quarts antifreeze, 4 quarts water

## Chapter 5 cumulative test

1. 1,756 2. undefined 3.  $-36$  4.  $5a+3b$   
 5.  $9x^2-12x+4$  6.  $25y^2-4$  7.  $4a^2-21ab-18b^2$   
 8.  $x^3-3x^2y+3xy^2-y^3$  9.  $-33$  10.  $\{-2\}$  11.  $\{8\}$   
 12.  $-\frac{9}{2} \leq x < 3$  13.  $x > 3$  14.  $(m+n)(x-y)$   
 15.  $(3a+4)(a+1)$  16.  $(2x-5)^2$  17.  $(2z+3)(2z-3)$   
 18.  $(6-y)(6+y)$  19.  $(x-15)(x+3)$   
 20.  $2(a-2b)(a^2+2ab+4b^2)$  21.  $\{7, -2\}$  22.  $\left\{\frac{3}{2}, -3\right\}$   
 23.  $\frac{z^3}{64y^3}$  24.  $a^3$  25.  $8a^3$  26.  $x=24$  27.  $\frac{13}{6}$  or 13:6  
 28.  $x-6+\frac{1}{x-2}$  29.  $\frac{4}{3}$  30.  $\frac{9b}{7a^2}$  31.  $\frac{a-6}{a-7}$   
 32.  $\frac{8}{5(x+y)}$  33.  $\frac{y+4}{y+5}$  34.  $\frac{x-3y}{3x+2y}$  35.  $21\frac{1}{3}$  inches  
 36.  $12\frac{6}{7}$  feet

## Chapter 6

## Exercise 6-1

## Answers to odd-numbered problems

1.  $\frac{3}{5}$  3.  $\frac{5}{6}$  5.  $2a$  7. 10 9.  $\frac{x}{4y}$  11.  $\frac{3x}{4}$  13.  $\frac{4}{35x}$   
 15.  $\frac{35x}{4}$  17.  $\frac{7c}{2b}$  19.  $\frac{x}{6y^2}$  21.  $\frac{16bcx}{3az}$  23.  $\frac{2a}{5b^2x}$   
 25.  $\frac{4}{x+y}$  27.  $-\frac{3}{4}$  29.  $-\frac{15}{4}$  31.  $\frac{4}{a-5}$  33.  $\frac{18(x-2)}{x+2}$   
 35.  $\frac{24y(x-2)}{25}$  37.  $(r-4)(r-1)$  39.  $-4(x+3)$   
 41.  $\frac{a-3}{a-5}$  43.  $\frac{(x-3)(x+1)}{(x-1)(x+2)}$  45.  $\frac{(2x-1)(x-1)}{(x-8)(x+7)}$   
 47. 1 49.  $3x+4$  51.  $\frac{1}{(2x+1)^2}$  53.  $\frac{2a+12}{3a^2+9a+27}$   
 55.  $\frac{(z-7)(z+3)}{5(z^2-2z+4)}$  57.  $y+5$

## Solutions to trial exercise problems

10.  $\frac{7a}{12b} \cdot \frac{9b}{28} = \frac{7a \cdot 9b}{12b \cdot 28} = \frac{7 \cdot a \cdot 3 \cdot 3 \cdot b}{2 \cdot 2 \cdot 3 \cdot b \cdot 2 \cdot 2 \cdot 7}$   
 $= \frac{3 \cdot a \cdot (7 \cdot 3 \cdot b)}{2 \cdot 2 \cdot 2 \cdot 2 \cdot (7 \cdot 3 \cdot b)} = \frac{3a}{16}$   
 20.  $\frac{28m}{15n} \div \frac{7m^2}{3n^3} = \frac{28m \cdot 3n^3}{15n \cdot 7m^2} = \frac{4n^2}{5m}$   
 21.  $\frac{24abc}{7xyz^2} \cdot \frac{14x^2yz}{9a^2} = \frac{3 \cdot 8 \cdot a \cdot b \cdot c \cdot 2 \cdot 7 \cdot x^2 \cdot y \cdot z}{7 \cdot x \cdot y \cdot z^2 \cdot 3 \cdot 3 \cdot a^2}$   
 $= \frac{8 \cdot b \cdot c \cdot 2 \cdot x(3 \cdot 7 \cdot a \cdot x \cdot y \cdot z)}{z \cdot 3 \cdot a(3 \cdot 7 \cdot a \cdot x \cdot y \cdot z)} = \frac{8 \cdot b \cdot c \cdot 2 \cdot x}{z \cdot 3 \cdot a} = \frac{16bcx}{3az}$   
 25.  $\frac{x+y}{3} \cdot \frac{12}{(x+y)^2} = \frac{2 \cdot 2 \cdot 3(x+y)}{3(x+y)^2} = \frac{4}{x+y}$   
 30.  $\frac{8y+16}{3-y} \cdot \frac{4y-12}{3y+6} = \frac{8(y+2) \cdot 4(y-3)}{-(y-3) \cdot 3(y+2)} = \frac{8 \cdot 4}{-3} = -\frac{32}{3}$   
 37.  $\frac{r^2-16}{r+1} \div \frac{r+4}{r^2-1} = \frac{(r^2-16)(r^2-1)}{(r+1)(r+4)}$   
 $= \frac{(r-4)(r+4)(r+1)(r-1)}{(r+1)(r+4)} = \frac{(r-4)(r-1)}{1}$   
 $= r^2 - 5r + 4$   
 39.  $\frac{9-x^2}{x+y} \cdot \frac{4x+4y}{x-3} = \frac{(3-x)(3+x) \cdot 4(x+y)}{(x+y)(x-3)}$   
 $= \frac{-1(x-3)(x+3) \cdot 4(x+y)}{(x+y)(x-3)} = \frac{-1(x+3) \cdot 4}{1}$   
 $= -4(x+3) = -4x-12$   
 41.  $\frac{a^2-5a+6}{a^2-9a+20} \cdot \frac{a^2-5a+4}{a^2-3a+2}$   
 $= \frac{(a-3)(a-2) \cdot (a-4)(a-1)}{(a-4)(a-5) \cdot (a-2)(a-1)} = \frac{a-3}{a-5}$   
 47.  $\frac{6r^2-r-7}{12r^2+16r-35} \div \frac{r^2-r-2}{2r^2+r-10}$   
 $= \frac{(6r^2-r-7)(2r^2+r-10)}{(12r^2+16r-35)(r^2-r-2)}$   
 $= \frac{(6r-7)(r+1)(2r+5)(r-2)}{(6r-7)(2r+5)(r-2)(r+1)} = 1$   
 49.  $(3x^2-2x-8) \div \frac{x^2-4}{x+2} = \frac{(3x+4)(x-2)}{1}$   
 $\cdot \frac{x+2}{(x+2)(x-2)} = 3x+4$

53.  $\frac{10}{a^3-27} \cdot \frac{a^2+3a-18}{15} = \frac{2 \cdot 5(a+6)(a-3)}{3 \cdot 5(a-3)(a^2+3a+9)}$   
 $= \frac{2(a+6)}{3(a^2+3a+9)} = \frac{2a+12}{3a^2+9a+27}$

## Review exercises

1.  $\frac{19}{12}$  2.  $\frac{11}{24}$  3.  $2(x+5)(x-5)$  4.  $(x+11)(x-2)$   
 5.  $(x+4)^2$  6.  $x=\frac{24}{5}$  7.  $y=15$  8.  $7.89 \times 10^{-5}$



## Exercise 6-2

## Answers to odd-numbered problems

1.  $\frac{8}{x}$  3.  $\frac{7}{p}$  5.  $\frac{14x}{x+2}$  7.  $-\frac{2}{x}$  9.  $\frac{x+2}{x^2-1}$   
 11.  $-\frac{1}{7}$  13.  $\frac{9}{z}$  15.  $\frac{-7}{x-2}$  17.  $\frac{9y}{y-6}$   
 19.  $\frac{-x+4}{x-5}$  21.  $\frac{3y+2}{2y-3}$  23.  $\frac{-x+4}{2x-5}$  25.  $18x$   
 27.  $48x^2$  29.  $140y^3$  31.  $64a^4$  33.  $180a^3$  35.  $3(x-4)$   
 37.  $18y^3(y-4)$  39.  $a(a+1)(a-1)$  41.  $8(a+2)(a+1)$   
 43.  $(a+2)(a-2)(a-3)$  45.  $(a+3)(a-3)(a-2)$   
 47.  $2(x-7)(x+7)$

## Solutions to trial exercise problems

5.  $\frac{5x}{x+2} + \frac{9x}{x+2} = \frac{5x+9x}{x+2} = \frac{14x}{x+2}$  8.  $\frac{3y-2}{y^2} - \frac{4y-1}{y^2}$   
 $= \frac{(3y-2)-(4y-1)}{y^2} = \frac{3y-2-4y+1}{y^2} = \frac{-y-1}{y^2}$   
 16.  $\frac{1}{x-7} - \frac{5}{7-x} = \frac{1}{x-7} + \frac{5}{x-7} = \frac{6}{x-7}$   
 21.  $\frac{2y-5}{2y-3} - \frac{y+7}{3-2y} = \frac{2y-5}{2y-3} + \frac{y+7}{2y-3} = \frac{3y+2}{2y-3}$   
 32.  $4x^2 = 2^2 \cdot x^2$   
 $3x = 3 \cdot x$   
 $8x^3 = 2^3 \cdot x^3$   
 LCD =  $2^3 \cdot 3 \cdot x^3 = 24x^3$   
 44.  $y^2 - y - 12 = (y-4)(y+3)$   
 $y^2 + 6y + 9 = (y+3)^2$   
 LCD =  $(y+3)^2(y-4)$   
 47.  $x^2 - 49 = (x+7)(x-7)$   
 $7-x = -1(x-7)$   
 $2x+14 = 2(x+7)$   
 LCD =  $2(x+7)(x-7)$

## Review exercises

1. commutative property of addition 2.  $5(y+2)(y-2)$   
 3.  $(x+10)^2$  4.  $(3y-4)(y+1)$  5.  $\left\{\frac{8}{19}\right\}$  6.  $\{-3, 5\}$

## Exercise 6-3

## Answers to odd-numbered problems

1.  $\frac{2x+9}{12}$  3.  $\frac{8x+1}{48}$  5.  $\frac{11}{3}$  7.  $\frac{15y-12}{(y+4)(y-5)}$  9.  $\frac{10}{y-2}$   
 11.  $\frac{7x+62}{4(x+2)(x-2)}$  13.  $\frac{9x+40}{x+8}$  15.  $\frac{x(x+4)}{(x+1)(x-1)}$   
 17.  $\frac{9x+22}{(x+3)(x-3)(x+2)}$  19.  $\frac{7y^2-13y}{(y-3)^2(y+1)}$   
 21.  $\frac{2y^2-9y+1}{(y-5)(y-3)(y+2)}$  23.  $\frac{2y-15}{18}$  25.  $\frac{25}{42y}$   
 27.  $\frac{19a+39}{60}$  29.  $\frac{4x+25}{18x}$  31.  $\frac{-5x-17}{(2x-3)(x-5)}$   
 33.  $\frac{17}{7(y+2)}$  35.  $\frac{9x+66}{x+8}$  37.  $\frac{-25x-9}{3x+1}$   
 39.  $\frac{-6a+3}{(a-2)(a+2)(a-3)}$  41.  $\frac{12y-12}{(y-6)(y+4)(y-3)}$   
 43.  $\frac{-3a^2+11a-3}{(a-3)(a-2)}$  45.  $\frac{13}{12z}$  47.  $\frac{52a}{45}$  49.  $\frac{39b-5}{36}$

51.  $\frac{-z^3+9z^2-13}{(z-2)(z+2)(z-1)(z+1)}$  53.  $\frac{bc+ac+ab}{abc}$  55.  $\frac{3}{m}$   
 57.  $\frac{h}{36}$  59.  $\frac{48}{m}$  61.  $w = \frac{A}{23}$  63.  $b = \frac{2A}{9}$  65.  $t = \frac{d}{55}$   
 67.  $\frac{b}{a}$

## Solutions to trial exercise problems

6.  $\frac{4}{x-1} + \frac{5}{x+3} = \frac{4(x+3)+5(x-1)}{(x-1)(x+3)}$   
 $= \frac{4x+12+5x-5}{(x-1)(x+3)} = \frac{9x+7}{(x-1)(x+3)}$   
 11.  $\frac{12}{x^2-4} + \frac{7}{4x-8} = \frac{12}{(x+2)(x-2)} + \frac{7}{4(x-2)}$   
 $= \frac{12 \cdot 4}{4(x+2)(x-2)} + \frac{7(x+2)}{4(x+2)(x-2)} = \frac{48+(7x+14)}{4(x+2)(x-2)}$   
 $= \frac{7x+62}{4(x+2)(x-2)}$  13.  $5 + \frac{4x}{x+8} = \frac{5(x+8)}{x+8} + \frac{4x}{x+8}$   
 $= \frac{5x+40+4x}{x+8} = \frac{9x+40}{x+8}$  25.  $\frac{9}{14y} - \frac{1}{21y} = \frac{9 \cdot 3}{2 \cdot 3 \cdot 7 \cdot y}$   
 $= \frac{1 \cdot 2}{2 \cdot 3 \cdot 7 \cdot y} = \frac{2}{42y} - \frac{2}{42y} = \frac{2-2}{42y} = \frac{0}{42y}$   
 27.  $\frac{5a+3}{12} - \frac{a-4}{10} = \frac{5(5a+3)}{60} - \frac{6(a-4)}{60}$   
 $= \frac{(25a+15)-(6a-24)}{60} = \frac{25a+15-6a+24}{60} = \frac{19a+39}{60}$   
 34.  $\frac{14}{5x-15} - \frac{8}{2x-6} = \frac{14}{5(x-3)} - \frac{8}{2(x-3)} = \frac{14 \cdot 2}{10(x-3)}$   
 $= \frac{8 \cdot 5}{10(x-3)} = \frac{28-40}{10(x-3)} = \frac{-12}{10(x-3)} = \frac{-6}{5(x-3)}$   
 42.  $\frac{2p}{p^2-9p+20} - \frac{5p-2}{p-5} = \frac{2p}{(p-5)(p-4)} - \frac{5p-2}{(p-5)(p-4)}$   
 $= \frac{2p-(5p-2)}{(p-5)(p-4)} = \frac{2p-5p+2}{(p-5)(p-4)} = \frac{-3p+2}{(p-5)(p-4)}$   
 52.  $\frac{1}{f} = (u-1) \left( \frac{1}{R_1} + \frac{1}{R_2} \right) = (u-1) \left( \frac{R_2}{R_1 R_2} + \frac{R_1}{R_1 R_2} \right)$   
 $= (u-1) \left( \frac{R_2+R_1}{R_1 R_2} \right)$  59. Since  $n$  is one of the numbers, let  $x$  be  
 the other number. Then  $x \cdot n = 48$  and  $x$  (the other number) =  $\frac{48}{n}$ .  
 62. Using  $A = \frac{1}{2}bh$ , then  $2A = bh$  (multiply each member by 2).  
 Then  $h = \frac{2A}{b} = \frac{2(21)}{b} = \frac{42}{b}$ .

## Review exercises

1.  $(x-7)^2$  2.  $(2x-1)(x-5)$  3.  $4(x+2)(x-2)$   
 4.  $x^2-81$  5.  $16x^2+24x+9$  6.  $2x^2-15x-8$   
 7. 48 8.  $12x^2$  9.  $(x-3)^2(x+3)$   
 10.  $\frac{13}{2x}$  11.  $\frac{2x+1}{(x-2)(x-1)}$

## Exercise 6-4

## Answers to odd-numbered problems

1.  $\frac{5}{6}$  3.  $\frac{3}{2}$  5.  $\frac{8}{9}$  7.  $\frac{5}{2}$  9.  $\frac{23}{30}$  11.  $\frac{11}{2}$   
 13.  $\frac{4x+1}{4x-3}$  15.  $\frac{3a+1}{2-4a}$  17.  $\frac{4a^2+3}{5a^2-3a}$  19.  $\frac{ab-3}{ab+4}$   
 21.  $\frac{y+x}{y-x}$  23.  $xy$  25.  $\frac{4b-5a^2}{a^3b-a^2b^2}$  27.  $-\frac{y}{x}$   
 29.  $\frac{2(x^2+y^2)}{(x^2-y^2)^2}$  31.  $\frac{2b+3}{a^3b^4}$  33.  $\frac{3a-1}{17-a}$  35.  $\frac{-5}{x+31}$   
 37.  $-\frac{3a^2+5a-8}{a+29}$  39.  $\frac{(y-3)(y+6)}{(y+1)(y-4)}$  41.  $\frac{T_1}{T_2-T_1}$   
 43.  $\frac{L_1L_2L_3}{L_2L_3+L_1L_3+L_1L_2}$

## Solutions to trial exercise problems

7.  $\frac{7}{2+\frac{4}{5}} = \frac{7 \cdot 5}{\left(2+\frac{4}{5}\right) \cdot 5} = \frac{7 \cdot 5}{2 \cdot 5 + \frac{4}{5} \cdot 5} = \frac{35}{10+4} = \frac{35}{14} = \frac{5}{2}$   
 12.  $\frac{\frac{6}{7} - \frac{5}{14}}{\frac{3}{14} - \frac{5}{7}} = \frac{\left(\frac{6}{7} - \frac{5}{14}\right) \cdot 14}{\left(\frac{3}{14} - \frac{5}{7}\right) \cdot 14} = \frac{\frac{6}{7} \cdot 14 - \frac{5}{14} \cdot 14}{\frac{3}{14} \cdot 14 - \frac{5}{7} \cdot 14} = \frac{12-5}{3-10} = \frac{7}{-7} = -1$   
 15.  $\frac{\frac{1}{a} + 3}{\frac{2}{a} - 4} = \frac{\left(\frac{1}{a} + 3\right) \cdot a}{\left(\frac{2}{a} - 4\right) \cdot a} = \frac{\frac{1}{a} \cdot a + 3 \cdot a}{\frac{2}{a} \cdot a - 4 \cdot a} = \frac{1+3a}{2-4a} = \frac{3a+1}{2-4a}$   
 22.  $\frac{\frac{3}{x^2} - \frac{4}{y}}{\frac{5}{x} + \frac{2}{y^2}} = \frac{\left(\frac{3}{x^2} - \frac{4}{y}\right) \cdot x^2y^2}{\left(\frac{5}{x} + \frac{2}{y^2}\right) \cdot x^2y^2} = \frac{\frac{3}{x^2} \cdot x^2y^2 - \frac{4}{y} \cdot x^2y^2}{\frac{5}{x} \cdot x^2y^2 + \frac{2}{y^2} \cdot x^2y^2} = \frac{3y^2 - 4x^2y}{5xy^2 + 2x^2}$   
 27.  $\frac{\frac{1}{x+y} - \frac{1}{x-y}}{\frac{1}{x+y} + \frac{1}{x-y}} = \frac{\left(\frac{1}{x+y} - \frac{1}{x-y}\right) \cdot (x+y)(x-y)}{\left(\frac{1}{x+y} + \frac{1}{x-y}\right) \cdot (x+y)(x-y)} = \frac{\frac{1}{x+y} \cdot (x+y)(x-y) - \frac{1}{x-y} \cdot (x+y)(x-y)}{\frac{1}{x+y} \cdot (x+y)(x-y) + \frac{1}{x-y} \cdot (x+y)(x-y)} = \frac{(x-y) - (x+y)}{(x-y) + (x+y)} = \frac{x-y-x-y}{x-y+x+y} = \frac{-2y}{2x} = \frac{-y}{x}$   
 36.  $\frac{\frac{7}{b-7} + \frac{8}{b-5}}{\frac{6}{b^2-12b+35}} = \frac{\left(\frac{7}{b-7} + \frac{8}{b-5}\right) \cdot (b-5)(b-7)}{\frac{6}{(b-5)(b-7)} \cdot (b-5)(b-7)}$   
 The LCD is  $(b-5)(b-7)$ .  

$$= \frac{\frac{7}{b-7} \cdot (b-5)(b-7) + \frac{8}{b-5} \cdot (b-5)(b-7)}{6} = \frac{7(b-5) + 8(b-7)}{6} = \frac{7b-35+8b-56}{6} = \frac{15b-91}{6}$$

## Review exercises

1.  $\frac{6}{7}$  2.  $\frac{x-7}{x+2}$  3.  $2x^3 + 3x^2 + x - 21$  4.  $\{3\}$  5.  $\{-2\}$   
 6.  $\{-3, 1\}$  7. Domain is all real numbers except  $-7$ . 8. Domain is all real numbers except  $-2$  and  $2$ . 9. a. undefined b.  $0$

## Exercise 6-5

## Answers to odd-numbered problems

1.  $\left\{\frac{8}{3}\right\}$  3.  $\left\{\frac{14}{3}\right\}$  5.  $\{-22\}$  7.  $\left\{\frac{10}{9}\right\}$  9.  $\left\{\frac{9}{2}\right\}$   
 11.  $\left\{-\frac{5}{8}\right\}, x \neq 0$  13.  $\left\{\frac{19}{24}\right\}, R \neq 0$  15.  $\left\{-\frac{5}{27}\right\}, y \neq 0$   
 17.  $\{1\}, b \neq 0$  19.  $\left\{\frac{4}{37}\right\}, p \neq 0$  21.  $\left\{-\frac{13}{2}\right\}, a \neq 0$   
 23.  $\{36\}, x \neq 4, -4$  25.  $\left\{\frac{47}{9}\right\}, z \neq \frac{-1}{3}$  27.  $\emptyset; y \neq 2$   
 29.  $\left\{\frac{11}{5}\right\}, b \neq 2, -2$  31.  $\{40\}, a \neq 5, -5$   
 33.  $\left\{-\frac{19}{3}\right\}, x \neq 2, -3, -5$  35.  $\left\{\frac{8}{3}\right\}, a \neq 3, -4$  37.  $\left\{-\frac{2}{3}\right\}$   
 39.  $\left\{\frac{3}{2}, -3\right\}$  41.  $\left\{\frac{2}{3}, -4\right\}$  43.  $\left\{\frac{2}{3}, -1\right\}$   
 45.  $x = \frac{2y}{3y-1}$  47.  $c_1 = \frac{cc_2}{c_2-c}$  49.  $a = \frac{8b}{b-8}$   
 51.  $a = \frac{3b-5}{4}$  53.  $x = \frac{-2(y+6)}{5}$  55.  $D = \frac{P}{pL}$   
 57.  $N = \frac{2P}{D-P}$  59.  $f = \frac{Fa}{A}$  61.  $T_2 = \frac{T_1P_2V_2}{P_1V_1}$   
 63.  $L_1 = \frac{L_2}{\alpha(t_2-t_1)+1}$  65.  $R_1 = \frac{RR_2}{R_2-R}$

## Solutions to trial exercise problems

2.  $\frac{4x}{5} - \frac{2}{3} = 4$  Multiply by 15 to get  
 The LCD is 15.  $\frac{4x}{5} \cdot 15 - \frac{2}{3} \cdot 15 = 4 \cdot 15$   
 $4x \cdot 3 - 2 \cdot 5 = 60$   
 $12x - 10 = 60$   
 $12x = 70$   
 $x = \frac{70}{12} = \frac{35}{6}$   
 The solution set is  $\left\{\frac{35}{6}\right\}$ .  
 6.  $\frac{3R}{4} - 5 = \frac{5}{6}$  Multiply by 12 to get  
 The LCD is 12.  $\frac{3R}{4} \cdot 12 - 5 \cdot 12 = \frac{5}{6} \cdot 12$   
 $3R \cdot 3 - 60 = 5 \cdot 2$   
 $9R - 60 = 10$   
 $9R = 70$   
 $R = \frac{70}{9}$   
 The solution set is  $\left\{\frac{70}{9}\right\}$ .



15.  $\frac{4}{6y} + 5 = \frac{1}{9y} + 2$  Multiply by  $18y$  to get

The LCD is  $18y$ .  $\frac{4}{6y} \cdot 18y + 5 \cdot 18y = \frac{1}{9y} \cdot 18y + 2 \cdot 18y$   
 $4 \cdot 3 + 90y = 1 \cdot 2 + 36y$   
 $12 + 90y = 2 + 36y$   
 $90y = -10 + 36y$   
 $54y = -10$   
 $y = \frac{-10}{54} = \frac{-5}{27}$

The solution set is  $\left\{-\frac{5}{27}\right\} (y \neq 0)$ .

22.  $\frac{R+2}{10R} + \frac{4R-1}{4R} = 2$  Multiply by  $20R$  to get

The LCD is  $20R$ .  $\frac{R+2}{10R} \cdot 20R + \frac{4R-1}{4R} \cdot 20R = 2 \cdot 20R$   
 $(R+2) \cdot 2 + (4R-1) \cdot 5 = 40R$   
 $2R + 4 + 20R - 5 = 40R$   
 $22R - 1 = 40R$   
 $-1 = 18R$   
 $R = \frac{-1}{18}$

The solution set is  $\left\{-\frac{1}{18}\right\} (R \neq 0)$ .

34.  $\frac{8}{a^2 - 6a + 8} = \frac{1}{a^2 - 16}$  Multiply both members by  $(a-2)(a+4)(a-4)$ .

$\frac{8}{(a-4)(a-2)} \cdot (a-2)(a+4)(a-4) = \frac{1}{(a+4)(a-4)} \cdot (a-2)(a+4)(a-4)$   
 $8(a+4) = 1(a-2)$   
 $8a + 32 = a - 2$   
 $8a = a - 34$   
 $7a = -34$   
 $a = \frac{-34}{7}$

The solution set is  $\left\{-\frac{34}{7}\right\} (a \neq 4, 2, -4)$ .

37.  $3x^2 + 4x + \frac{4}{3} = 0$  Multiply by 3 to obtain

$9x^2 + 12x + 4 = 0$  Factor the left member.  
 $(3x+2)^2 = 0$   
 $3x+2 = 0$   
 $3x = -2$   
 $x = -\frac{2}{3}$  So

The solution set is  $\left\{-\frac{2}{3}\right\}$ .

42.  $x^2 - \frac{5}{6}x = \frac{2}{3}$  Multiply by the LCD, 6, to obtain

$6x^2 - 5x = 4$   
 $6x^2 - 5x - 4 = 0$   
 $(2x+1)(3x-4) = 0$   
 $2x+1 = 0$  or  $3x-4 = 0$   
 $x = -\frac{1}{2}$  or  $x = \frac{4}{3}$

The solution set is  $\left\{-\frac{1}{2}, \frac{4}{3}\right\}$ .



47.  $\frac{1}{c} = \frac{1}{c_1} + \frac{1}{c_2}$  Multiply by  $cc_1c_2$ .

$$\frac{1}{c} \cdot cc_1c_2 = \frac{1}{c_1} \cdot cc_1c_2 + \frac{1}{c_2} \cdot cc_1c_2$$

$$c_1c_2 = cc_2 + cc_1 \quad \text{Add } -cc_1 \text{ to both members.}$$

$$c_1c_2 - cc_1 = cc_2 \quad \text{Factor } c_1 \text{ in the left member.}$$

$$(c_2 - c)c_1 = cc_2 \quad \text{Divide both members by } c_2 - c.$$

$$c_1 = \frac{cc_2}{c_2 - c}$$

52.  $\frac{5}{m} + 4 = \frac{6a}{2m} + 3b$  Multiply by the LCD  $2m$ .

$$\frac{5}{m} \cdot 2m + 4 \cdot 2m = \frac{6a}{2m} \cdot 2m + 3b \cdot 2m$$

$$10 + 8m = 6a + 6bm \quad \text{Add } -10 \text{ to both members.}$$

$$8m = 6a - 10 + 6bm \quad \text{Add } -6bm \text{ to both members.}$$

$$8m - 6bm = 6a - 10 \quad \text{Factor } m \text{ in the left member.}$$

$$(8 - 6b)m = 6a - 10 \quad \text{Divide both members by } 8 - 6b.$$

$$m = \frac{6a - 10}{8 - 6b} = \frac{2(3a - 5)}{2(4 - 3b)} = \frac{3a - 5}{4 - 3b}$$

58.  $\frac{T_A}{T_B} = \frac{R_B}{R_A}$  Multiply by the LCD  $T_BR_A$ .

$$\frac{T_A}{T_B} \cdot T_BR_A = \frac{R_B}{R_A} \cdot T_BR_A$$

$$T_AR_A = R_BT_B$$

$$T_B = \frac{T_AR_A}{R_B}$$

62. Given  $k = \frac{L_t - L_0}{L_0t}$ . Multiply both members by  $L_0t$  to get  $kL_0t = L_t - L_0$ .

a. To find  $L_t$ , add  $L_0$  to both members to get  $L_t = kL_0t + L_0$ .

b. To find  $L_0$ , use  $L_t = kL_0t + L_0$ , factor  $L_0$  in the right member to get  $L_t = (kt + 1)L_0$ . Divide both members by  $kt + 1$ .

$$\text{Then } \frac{L_t}{kt + 1} = L_0 \text{ or } L_0 = \frac{L_t}{kt + 1}.$$

### Review exercises

1. 14    2.  $\frac{17}{3}$     3. 4

4.  $y = -5x + 4$     5.  $y = \frac{-2x + 6}{3}$

6.  $y = \frac{x - 8}{4}$     7.  $\frac{3y}{x^2}$     8.  $\frac{y^4}{x^5}$

### Exercise 6-6

#### Answers to odd-numbered problems

1.  $29\frac{1}{6}$  minutes    3.  $2\frac{14}{29}$  hours    5. 12 hours    7.  $3\frac{1}{3}$  hours

9. 10 hours    11. 110 miles at 55 mph, 210 miles at 60 mph

13. car A, 40 mph; car B, 50 mph    15.  $\frac{6}{7}$  mph    17. 220 miles

19.  $\frac{5}{9}$     21. 3 and 12    23. 6    25. 4    27. adult, 15 pills; child,

9 pills    29. \$38.21    31.  $\frac{60}{17}$  ohms or  $3\frac{9}{17}$  ohms    33. 20 ohms

35. 60 ohms

## Solutions to trial exercise problems

1. Let  $x$  = the number of minutes required for both boys to mow the lawn. Then,

$$\frac{1}{50} + \frac{1}{70} = \frac{1}{x}$$

Multiply by the LCM of 50, 70, and  $x$ , that is,  $350x$ .

$$\begin{aligned} 350x \cdot \frac{1}{50} + 350x \cdot \frac{1}{70} &= 350x \cdot \frac{1}{x} \\ 7x + 5x &= 350 \\ 12x &= 350 \\ x &= \frac{350}{12} = 29\frac{1}{6} \end{aligned}$$

Together Jim and Kenny could mow the lawn in  $29\frac{1}{6}$  minutes.

5. Let  $x$  = the time required for Dick to paint the house alone.

$$\text{Then } \frac{1}{6} + \frac{1}{x} = \frac{1}{4}.$$

Multiply both members by the LCM of 4, 6, and  $x$ , that is,  $12x$ .

$$\begin{aligned} 12x \cdot \frac{1}{6} + 12x \cdot \frac{1}{x} &= 12x \cdot \frac{1}{4} \\ 2x + 12 &= 3x \\ 12 &= x \end{aligned}$$

Therefore Dick would take 12 hours to paint the house alone.

13. Let  $x$  = the average speed of car A.

Then  $x + 10$  = the average speed of car B.

Using  $t = \frac{d}{r}$ , let  $t_A$  = time of car A,  $t_B$  = time of car B,

then  $t_A = t_B$ .

Now  $t_A = \frac{120}{x}$  and  $t_B = \frac{150}{x + 10}$ , so

$$\frac{120}{x} = \frac{150}{x + 10}$$

Multiply both members by the LCM of  $x$  and  $x + 10$ , that is,  $x(x + 10)$ .

$$x(x + 10) \cdot \frac{120}{x} = x(x + 10) \cdot \frac{150}{x + 10}$$

$$(x + 10)120 = 150x$$

$$120x + 1,200 = 150x$$

$$1,200 = 30x$$

$$40 = x$$

So  $x + 10 = 50$ . Therefore car B averaged 50 mph and car A averaged 40 mph.

20. Let  $x$  = the numerator of the fraction. Then  $x + 7$  = the denominator of the fraction.

Then  $\frac{x + 3}{(x + 7) - 1} = \frac{4}{5}$ . Simplifying, we have the equation

$$\frac{x + 3}{x + 6} = \frac{4}{5}.$$

Multiply both members by the LCM of 5 and  $x + 6$ , that is,  $5(x + 6)$ .

$$5(x + 6) \cdot \frac{x + 3}{x + 6} = 5(x + 6) \cdot \frac{4}{5}$$

$$5(x + 3) = 4(x + 6)$$

$$5x + 15 = 4x + 24$$

$$x + 15 = 24$$

$$x = 9$$

$$x + 7 = 16$$

The original fraction is then  $\frac{9}{16}$ .

21. Let  $x$  = the lesser of the two numbers then

$4x$  = the greater of the two numbers.

Their reciprocals are then  $\frac{1}{x}$  and  $\frac{1}{4x}$  and we get the equation

$$\frac{1}{x} + \frac{1}{4x} = \frac{5}{12}.$$

Multiply both members by the LCM of  $x$ ,  $4x$ , and 12, that is,  $12x$ .

$$12x \cdot \frac{1}{x} + 12x \cdot \frac{1}{4x} = 12x \cdot \frac{5}{12}$$

$$12 + 3 = 5x$$

$$15 = 5x$$

$$x = 3$$

then

$$4x = 12$$

Therefore the two numbers are 3 and 12.

(Note: To check, show  $\frac{1}{3} + \frac{1}{12} = \frac{5}{12}$ )

$$\frac{4}{12} + \frac{1}{12} = \frac{5}{12}$$

$$\frac{5}{12} = \frac{5}{12}$$

28. Let  $x$  = hourly wage of a journeyman electrician, then

$\frac{3}{8}x$  = the hourly wage of an apprentice electrician.

$$x + \frac{3}{8}x = 29.70$$

$$8 \cdot x + 8 \cdot \frac{3}{8}x = 8 \cdot 29.70$$

$$8x + 3x = 237.60$$

$$11x = 237.60$$

$$x = \$21.60$$

$$\frac{3}{8}x = \$8.10$$

The journeyman earns \$21.60 per hour and the apprentice earns \$8.10 per hour.

30. Let  $R$  = the total resistance of the circuit. Then

$$\frac{1}{6} + \frac{1}{8} = \frac{1}{R}.$$

Multiply by the LCM of 6, 8, and  $R$ , that is,  $24R$ .

$$24R \cdot \frac{1}{6} + 24R \cdot \frac{1}{8} = 24R \cdot \frac{1}{R}$$

$$4R + 3R = 24$$

$$7R = 24$$

$$R = \frac{24}{7}$$

Therefore the total resistance in the parallel circuit is  $\frac{24}{7}$

or  $3\frac{3}{7}$  ohms.

33. Let  $R_1$  = the resistance in the unknown branch. Then

$$\frac{1}{R_1} + \frac{1}{30} = \frac{1}{12}.$$

Multiply by the LCM of  $R_1$ , 12, and 30, that is,  $60R_1$ .

$$60R_1 \cdot \frac{1}{R_1} + 60R_1 \cdot \frac{1}{30} = 60R_1 \cdot \frac{1}{12}$$

$$60 + 2R_1 = 5R_1$$

$$60 = 3R_1$$

$$20 = R_1$$

Therefore the other branch has resistance of 20 ohms.

## Review exercises

1.  $\{2\}$  2.  $y = \frac{3x+6}{2}$  3.  $8(y+2)(y-2)$   
 4.  $(x+10)^2$  5.  $(3y+2)(y-2)$  6.  $\frac{5x^2+7x}{(x-1)(x+3)}$   
 7.  $\frac{-2y^2-23y}{(2y+1)(y-5)}$

## Chapter 6 review

1.  $\frac{9}{2}$  2.  $\frac{9a}{b}$  3.  $\frac{2x-y}{2x+y}$  4.  $x-2$  5.  $\frac{x(m-n)}{2}$   
 6.  $\frac{1}{y^2-2y+1}$  7. 2 8.  $-\frac{1}{3x+15}$  9.  $\frac{x-1}{x+1}$   
 10.  $\frac{2a}{3}$  11.  $\frac{8b}{21}$  12.  $\frac{9}{2ab}$  13.  $\frac{14x-7}{9x+12}$   
 14.  $\frac{1}{x^2+4x-12}$  15.  $\frac{10}{3}$  16.  $\frac{1}{x-8}$  17.  $\frac{3-b}{b^2+2b-3}$   
 18.  $\frac{3a+3}{4a-4}$  19.  $36x^2$  20.  $42a^2b^2$  21.  $2(x+3)(x-3)$   
 22.  $(y+5)(y-5)(y+3)$  23.  $4(z+2)(z-2)(z+1)$   
 24.  $x(x+1)^2(3x-5)$  25.  $\frac{3}{x}$  26.  $\frac{4y}{y-2}$  27.  $\frac{4x+4}{3x-1}$   
 28.  $\frac{-x+13}{2x-5}$  29.  $\frac{68x}{105}$  30.  $\frac{23}{48a}$   
 31.  $\frac{47x-6}{(3x+1)(4x-3)}$  32.  $\frac{4x^2-23x}{(x+4)(x-4)}$   
 33.  $\frac{80a+48b-15ab}{20a^2b^2}$  34.  $\frac{4a^2+4a+6}{a+1}$  35.  $\frac{-10x^2-3}{x^2+1}$   
 36.  $\frac{16x^2+5x+15}{24x^2}$  37.  $\frac{13y^2-89y}{(y-9)(y+2)(y-2)}$   
 38.  $\frac{2-2ax-36a}{(x+3)(x-3)}$  39.  $\frac{8x^2-40x-180}{x(x-5)(x+4)}$   
 40.  $\frac{10x-5y}{(x+y)^2(x-2y)}$  41.  $\frac{16}{63}$  42.  $\frac{1}{7}$  43. 1 44.  $\frac{y-x}{y+x}$   
 45.  $\frac{4+3x}{2-5x}$  46.  $\frac{a^2-b^2}{a^2+b^2}$  47.  $y-x$  48.  $\frac{xy}{x+y}$   
 49.  $\frac{a^3-2a^2b^2-3ab^2}{ab-3b^2}$  50.  $\{-96\}$  51.  $\left\{\frac{41}{48}\right\}$  52.  $\left\{\frac{13}{24}\right\}$   
 53.  $\left\{\frac{6}{5}\right\}$  54.  $\left\{-\frac{39}{14}\right\}$  55.  $\left\{\frac{7}{52}\right\}$  56.  $\{0,12\}$   
 57.  $\left\{-\frac{3}{2}, \frac{3}{2}\right\}$  58.  $\{-5,3\}$  59.  $\left\{6, \frac{3}{2}\right\}$  60.  $x = \frac{a+b}{3}$   
 61.  $x = \frac{y+3}{1-y}$  62.  $y = \frac{4a-3b}{a}$  63.  $y = 4b+5c$   
 64.  $l = \frac{Aey}{F}$  65.  $L = \frac{Wp-2E}{EF}$  66.  $R = \frac{Mr}{M-2}$   
 67.  $m = \frac{Fr}{v^2+gr}$  68.  $5\frac{1}{3}$  hr, 16 hr 69.  $1\frac{5}{7}$  mph 70.  $\frac{2}{3}$  or 1

## Chapter 6 cumulative test

1.  $-\frac{71}{12}$  2.  $\frac{3}{4}$  3.  $\frac{43}{30}$  4.  $\frac{8}{5}$  5.  $x^{11}$  6.  $\frac{1}{y^9}$  7.  $\frac{25x^4}{y^6}$   
 8.  $10x^3-4x^2+11x-9$  9.  $\left\{-\frac{26}{7}\right\}$  10.  $\left\{\frac{81}{4}\right\}$   
 11.  $\left\{-\frac{3}{2}, 2\right\}$  12.  $x \geq -5$  13.  $2(x+3)(x-3)$   
 14.  $3x^2(2x^3-12x+3)$  15.  $(2x+5)(2x+3)$   
 16.  $3(1+2x^3)(1-2x^3)$  17.  $49x^2-84x+36$   
 18.  $25-\frac{1}{4}x^2$  19.  $6x^3-19x^2+37x-33$   
 20.  $15-22x^2+8x^4$  21.  $x = \frac{120}{7}$   
 22. 2,400 foot-pounds/min 23.  $\frac{4a^2cf\ell}{5b^2}$  24.  $\frac{y^2+10y+21}{y^2-8y+12}$   
 25.  $y^2-5y+6$  26.  $\frac{(x-y)(x-1)}{y}$  27.  $\frac{2y+4x}{x^2y^2}$   
 28.  $\frac{3x+9y}{(x-y)(x+y)}$  29.  $\frac{9}{a-2}$  30.  $\frac{13x+56}{(x-7)(x+7)(x+2)}$   
 31.  $\frac{-x^2-6x}{(x+3)(x-2)(x-6)}$  32.  $\frac{3y-1}{4y+5}$  33.  $\frac{y+x}{3x-4y}$   
 34.  $\frac{x^2-11x+28}{x-5}$  35.  $\{2\}$  36.  $\left\{-\frac{1}{3}, 6\right\}$  37.  $q = \frac{fp}{p-f}$   
 38.  $2\frac{2}{5}$  days 39.  $\frac{8}{16}$

## Chapter 7

## Exercise 7-1

## Answers to odd-numbered problems

1. (1,2) and  $(-1,-4)$  are solutions. 3.  $(-1,2)$ ,  $(3,0)$  are solutions.  
 5. (1,2) is a solution. 7.  $(2,3)$ ,  $(0,0)$  are solutions.  
 9.  $(-4,1)$  and  $(-4,-4)$  are solutions. 11.  $(-5,3)$ ,  $(-5,8)$  are solutions.  
 13.  $(1,5)$ ,  $(-2,-4)$ ,  $(0,2)$  15.  $(3,-3)$ ,  $(-4,11)$ ,  $(0,3)$   
 17.  $(3,-5)$ ,  $(-2,10)$ ,  $(0,4)$  19.  $(-2,-1)$ ,  $(3,0)$ ,  $\left(0,-\frac{3}{5}\right)$   
 21.  $(1,-4)$ ,  $(-1,1)$ ,  $\left(0,-\frac{3}{2}\right)$  23.  $(1,5)$ ,  $(-6,5)$ ,  $(0,5)$   
 25.  $(7,-1)$ ,  $\left(-\frac{3}{5}, -1\right)$ ,  $(0,-1)$  27.  $(5,1)$ ,  $(-1,-2)$ ,  $(3,0)$   
 29.  $(3,2)$ ,  $(-6,-4)$ ,  $(0,0)$  31.  $(1,-2)$ ,  $(1,7)$ ,  $(1,0)$   
 33. a. (75,\$170) b. (300,\$620) c. (1,000,\$2,020) 35. a. (2,236)  
 b. (12,216) c. (0,240) 37. a. (0,20) b. (5,35) c. (3,29)  
 39. a. (3,165) b. (8,440) c.  $\left(\frac{26}{5}, 286\right)$



# Campfire queen Cycling champion Sentimental geologist\*

Learn more about  
Marjon Walrod  
and tell us more  
about you. Visit  
[pwc.com/bringit](http://pwc.com/bringit).

Your life. You can  
bring it with you.



\*connectedthinking

PRICEWATERHOUSECOOPERS 

© 2006 PricewaterhouseCoopers LLP. All rights reserved. "PricewaterhouseCoopers" refers to PricewaterhouseCoopers LLP (a Delaware limited liability partnership) or, as the context requires, the PricewaterhouseCoopers global network or other member firms of the network, each of which is a separate and independent legal entity. \*connectedthinking is a trademark of PricewaterhouseCoopers LLP (US). We are proud to be an Affirmative Action and Equal Opportunity Employer.

# Contents

20 Point Learning System	xi
Preface	xvii
Study Tips	xxiii

## Chapter 1 ■ Operations with real numbers



1-1 Operations with fractions	2
1-2 Operations with decimals and percents	14
1-3 The set of real numbers and the real number line	25
1-4 Addition of real numbers	34
1-5 Subtraction of real numbers	41
1-6 Multiplication of real numbers	46
1-7 Division of real numbers	51
1-8 Properties of real numbers and order of operations	56
Chapter 1 lead-in problem	62
Chapter 1 summary	63
Chapter 1 error analysis	64
Chapter 1 critical thinking	64
Chapter 1 review	64

## Chapter 2 ■ Solving equations and inequalities




2-1 Algebraic notation and terminology	67
2-2 Evaluating algebraic expressions	72
2-3 Algebraic addition and subtraction	79
2-4 The addition and subtraction property of equality	86
2-5 The multiplication and division property of equality	93
2-6 Solving linear equations	98
2-7 Solving literal equations and formulas	104
2-8 Word problems	107
2-9 Solving linear inequalities	113
Chapter 2 lead-in problem	122
Chapter 2 summary	122
Chapter 2 error analysis	123
Chapter 2 critical thinking	123
Chapter 2 review	124
Chapter 2 cumulative test	125



**Chapter 3 ■ Polynomials and exponents**

3-1	Exponents—I	127
3-2	Products of algebraic expressions	133
3-3	Exponents—II	139
3-4	Exponents—III	145
3-5	Scientific notation	148
	Chapter 3 lead-in problem	151
	Chapter 3 summary	152
	Chapter 3 error analysis	152
	Chapter 3 critical thinking	152
	Chapter 3 review	153
	Chapter 3 cumulative test	154

**Chapter 4 ■ Factoring and solution of quadratic equations by factoring**

4-1	Common factors	155
4-2	Factoring trinomials of the form $x^2 + bx + c$	162
4-3	Factoring trinomials of the form $ax^2 + bx + c$	166
4-4	Factoring the difference of two squares and perfect square trinomials	175
4-5	Other types of factoring	179
4-6	Factoring: A general strategy	184
4-7	Solving quadratic equations by factoring	186
4-8	Applications of the quadratic equation	193
	Chapter 4 lead-in problem	197
	Chapter 4 summary	198
	Chapter 4 error analysis	198
	Chapter 4 critical thinking	198
	Chapter 4 review	199
	Chapter 4 cumulative test	200

**Chapter 5 ■ Rational Expressions, Ratio and Proportion**

5-1	Rational numbers and rational expressions	202
5-2	Simplifying rational expressions	207
5-3	The quotient of two polynomials	212
5-4	Ratio and proportion	219
	Chapter 5 lead-in problem	228
	Chapter 5 summary	228
	Chapter 5 error analysis	228
	Chapter 5 critical thinking	229
	Chapter 5 review	229
	Chapter 5 cumulative test	230



## Chapter 6 ■ Operations with Rational Expressions



6-1	Multiplication and division of rational expressions	232
6-2	Addition and subtraction of rational expressions	239
6-3	Addition and subtraction of rational expressions	245
6-4	Complex fractions	253
6-5	Rational equations	258
6-6	Rational expression applications	265
	Chapter 6 lead-in problem	270
	Chapter 6 summary	271
	Chapter 6 error analysis	271
	Chapter 6 critical thinking	272
	Chapter 6 review	272
	Chapter 6 cumulative test	274

## Chapter 7 ■ Linear Equations in Two Variables




7-1	Ordered pairs and the rectangular coordinate system	276
7-2	Graphs of linear equations	289
7-3	The slope of a line	297
7-4	The equation of a line	305
7-5	Graphing linear inequalities in two variables	315
7-6	Functions defined by linear equations in two variables	323
	Chapter 7 lead-in problem	329
	Chapter 7 summary	329
	Chapter 7 error analysis	330
	Chapter 7 critical thinking	330
	Chapter 7 review	331
	Chapter 7 cumulative test	333

## Chapter 8 ■ Systems of Linear Equations




8-1	Solutions of systems of linear equations by graphing	335
8-2	Solutions of systems of linear equations by elimination	340
8-3	Solutions of systems of linear equations by substitution	346
8-4	Applications of systems of linear equations	351
8-5	Solving systems of linear inequalities by graphing	360
	Chapter 8 lead-in problem	363
	Chapter 8 summary	364
	Chapter 8 error analysis	364
	Chapter 8 critical thinking	364
	Chapter 8 review	365
	Chapter 8 cumulative test	366

## Chapter 9 ■ Roots and Radicals



9-1	Principal roots	367
9-2	Product property for radicals	373
9-3	Quotient property for radicals	377
9-4	Sums and differences of radicals	383
9-5	Further operations with radicals	386
9-6	Fractional exponents	391
9-7	Equations involving radicals	395
	Chapter 9 lead-in problem	400
	Chapter 9 summary	400
	Chapter 9 error analysis	401
	Chapter 9 critical thinking	401
	Chapter 9 review	401
	Chapter 9 cumulative test	402

## Chapter 10 ■ Solutions of Quadratic Equations



10-1	Solutions of quadratic equations by extracting the roots	404
10-2	Solutions of quadratic equations by completing the square	409
10-3	Solutions of quadratic equations by the quadratic formula	416
10-4	Complex solutions to quadratic equations	424
10-5	The graphs of quadratic equations in two variables—quadratic functions	432
	Chapter 10 lead-in problem	444
	Chapter 10 summary	445
	Chapter 10 error analysis	445
	Chapter 10 critical thinking	446
	Chapter 10 review	446
	Final examination	447
	Appendix Answers and Solutions	449
	Index	505



# Index

## A

Abscissa of a point, 281  
 Absolute value, 31  
 Addition  
   of decimals, 16  
   of fractions, 7–9, 239  
   of rational expressions, 239, 246  
 Addition, identity element of, 35  
 Addition, properties of  
   associative, 38  
   commutative, 35  
   identity, 35  
   rational expressions, 239, 246  
 Addition and subtraction property of equality, 88, 340  
 Addition and subtraction property of inequalities, 115  
 Addition of  
   algebraic expressions, 80–81  
   like radicals, 384–85  
   more than two real numbers, 42  
   rational expressions, 239, 246  
   two negative numbers, 36  
   two numbers with different signs, 36–37  
   two positive numbers, 34  
   two real numbers, 38  
 Additive inverse, 37–38  
 Algebraic expression, 67  
 Approximately equal to, 19, 29, 369  
 Associative property of  
   addition, 38  
   multiplication, 49  
 Axes, 280  
 Axis of symmetry, 437

## B

Base, 56, 127  
 Binomial, 68  
 Binomial, square of, 135, 410  
 Boundary line, 316  
 Braces, 25, 42  
 Brackets, 42

## C

Cantor, Georg, 25  
 Coefficient, 68  
 Common denominator, 7  
 Commutative property of  
   addition, 35  
   multiplication, 46  
 Completely factored form, 157

Completing the square, 411–13  
 Complex fractions, 253  
   primary denominator, 253  
   primary numerator, 253  
   secondary denominators of, 253  
   simplifying, 253–55  
 Complex numbers, 425  
   addition of, 425  
   multiplication of, 426  
   rationalizing the denominator of, 427  
 Components of an ordered pair, 277  
 Compound inequality, 114  
 Conjugate factors, 388  
 Consistent system of equations, 337  
 Constant, 67  
 Coordinate, 29, 281  
 Coordinates of a point, 281  
 Counting numbers, 26

## D

Decimal number, 14  
   fraction, 14  
   point, 14  
   repeating, 19, 27  
   terminating, 27  
 Decreasing, 29  
 Degree of a polynomial, 69, 186  
 Denominator  
   of a fraction, 2  
   least common, 42  
   of a rational expression, 203  
 Denominator, rationalizing a, 427, 480, 482  
 Dependent system, 337  
 Descending powers, 69, 163  
 Difference of two squares, 136  
 Discriminant, 429  
 Distance-rate-time, 266, 352  
 Distributive property of  
   multiplication over addition, 79  
 Dividend, 51  
 Division  
   monomial by monomial, 212  
   polynomial by monomial, 212  
   polynomial by polynomial, 213–16  
   property of rational expressions, 234  
   of rational expressions, 235  
 Division, definition of, 51  
 Division by zero, 53  
 Division involving zero, 53  
 Division of  
   decimals, 17  
   fractions, 5, 235  
   like bases, 140  
   two or more real numbers, 52  
   two real numbers, 52

Divisor, 51  
 Domain  
   of a rational expression, 204–5  
   of a relation, 323

## E

Element of a set, 25  
 Elimination, method of solving systems of linear equations, 340–44  
 Empty set, 260, 398  
 Equality, properties of  
   addition and subtraction, 88  
   multiplication and division, 94  
   squaring, 396  
   symmetric, 89  
 Equation, parts of an, 86  
 Equation of a line, 305  
 Equations  
   conditional, 87  
   diagram of, 86  
   equivalent, 87  
   first-degree, 87  
   graph of, 282–83, 289–94  
   identical, 87  
   linear, 87, 276  
   literal, 104  
   quadratic, 186, 404–31  
   radical, 395–96  
   rational, 258  
   system of, 335  
 Equivalent rational expressions, 245–46  
 Evaluating rational expressions, 203–4  
 Evaluation, 72  
 Expanded form, 56, 127  
 Exponent, 56, 127  
 Exponential form, 56, 127  
 Exponents, properties and definitions of  
   definition, 127  
   fraction, 139  
   fractional, 392  
   group of factors to a power, 129  
   negative exponents, 141  
   power of a power, 130  
   product, 128  
   quotient, 140  
   zero as an exponent, 142  
 Extracting the roots, 405–6  
 Extraneous solutions, 260, 396

## F

Factor, 3, 46  
 Factor, greatest common, 156  
 Factored form, completely, 157



**Factoring**

- common factors, 156
- difference of two cubes, 179–80
- difference of two squares 175–76
- four-term polynomials, 159
- by inspection, 170–74
- perfect square trinomials, 177–78
- strategy, 184
- sum of two cubes, 181–82
- trinomials, 162–64, 166–74

**FOIL, 135****Formulas, 74, 104****Four-term polynomials, 159****Fraction, 2**

- complex, 253
- improper, 2
- proper, 2

**Fraction exponents, 392****Fraction to a power, 139****Function, 324**

- domain of a, 325
- range of a, 325

**Fundamental principle of rational expressions, 207****G****Graph**

- of a linear equation, in two variables, 282–83, 289–94
- of a point, 29, 359
- a quadratic equation, 432–40
- of systems of equations, 336–37

**Graphing**

- a linear equation in two variables, 289–94
- linear inequalities in two variables, 315–19
- quadratic functions, 434–39
- systems of linear equations, 336–37
- systems of linear inequalities, 360–61

**Greatest common factor, 155–56****Grouping symbols, 42, 82****Grouping symbols, removing, 82****Group of factors to a power, 129****H****Half-planes, 316****Horizontal line, 292–93**

- slope of, 300

**I****Identical equation, 87****Identity, 87****Identity element**

- of addition, 35
- of multiplication, 47

**Inconsistent system, 337****Increasing, 29****Independent system, 337****Indeterminate, 53****Index, 370****Inequalities**

- compound, 114

**linear, 113**

- strict, 30
- in two variables, 315–19
- weak, 30

**Inequalities, properties of**

- addition and subtraction, 115
- multiplication and division, 115

**Inequality symbols, 30, 113****Integers, 26****Intercepts, x- and y-, 290****Inverse property**

- additive, 38
- multiplicative, 94

**Irrational numbers, 27, 368****L****Least common denominator (LCD), 7, 242****Least common multiple (LCM), 242****Like bases, 128****Like radicals, 383****Like terms, 80****Linear equation, 87****Linear equations in two variables, 276**

- graphs of, 289–93
- systems of, 335

**Linear inequalities in two variables, 315**

- graphs of, 315–19

**Linear inequality, 113****Lines**

- equation of, 305
- horizontal, 300
- parallel, 309
- perpendicular, 310
- point-slope form, 306
- slope-intercept form of, 307
- slope of, 297–302
- standard form of, 305
- vertical, 301

**Literal equation, 104****M****Mathematical statement, 86****Member of an equation, 86****Member of a set, 25****Mixed number, 5****Monomial, 68****Multinomial, 69****Multiple, least common, 242****Multiplication**

- of decimals, 16
- of fractions, 4
- identity element of, 47
- of rational expressions, 233
- symbols for, 46

**Multiplication, properties of**

- associative, 49
- commutative, 46
- fractions, 4, 232
- identity, 47

**rational expressions, 233****Multiplication and division property for inequalities, 115****Multiplication and division property of equality, 94****Multiplication of**

- like bases, 128
- monomial and a multinomial, 133
- monomials, 130
- multinomials, 134
- $n$ th roots, 374
- square roots, 373
- two negative numbers, 47
- two numbers with different signs, 47
- two or more real numbers, 48
- two positive numbers, 46
- two real numbers, 48

**Multiplicative inverse, 94****N****Natural numbers, 26****Negative exponents, 141****Negative integers, 26****Negative of, 31** **$n$ th root**

- addition of, 383–85
- definition of, 370
- of a fraction, 378, 380
- rationalizing, 379–81, 388

**Number line, 29****Numerator, 2, 203****Numerical coefficient, 68****O****Open sentence, 86****Opposite of, 31****Ordered pairs, first component of, 277****Ordered pairs, second component of, 277****Ordered pairs of numbers, 277****Order of operations, 57****Order relationships, 30****Ordinate of a point, 281****Origin, 29, 280****of a rectangular coordinate plane, 280****P****Parabola, 434**

- axis of symmetry of a, 437
- graphing a, 434–39
- intercepts of a, 435
- vertex of a, 436–37

**Parallel lines, 309****Parentheses, 42****Percent, 19–20****Percentage, 20–21****Perfect cube, 180****Perfect square, 175–76****Perfect square integer, 368****Perfect square trinomial, 135, 409****Perimeter, 10, 74****Perpendicular lines, 310****Pi ( $\pi$ ), 28****Point**

- abscissa of a, 281
- graph of a, 281
- ordinate of a, 281

**Point-slope form, 306****Polynomial, 68**

Polynomial, degree of, 69, 186  
 Positive integers, 26  
 Power of a power, 130  
 Prime factor form, 3  
 Prime number, 3  
 Prime polynomial, 164  
 Principal  $n$ th root, 370  
 Principal square root, 368  
 Product, 3, 46  
 Product of rational expressions, 233  
 Product property of exponents, 128  
 Product property of radicals, 374  
 Product property of square roots, 373  
 Properties of real numbers, 56  
 Proportion, 221  
   property of, 221  
   terms of, 221  
 Pythagorean Theorem, 369

**Q**

Quadrants, 280  
 Quadratic  
   equations, 186, 404–31  
   formula, 417–19  
   standard form of, 187, 404  
 Quotient, 5, 51  
   of two polynomials, 212–16  
 Quotient property of exponents, 140  
 Quotient property of radicals, 378, 380

**R**

Radical equation, 395–96  
 Radicals, properties of  
   product, 373, 374  
   quotient, 378, 380  
 Radical symbol, 368  
 Radicand, 368  
 Range of a relation, 323  
 Ratio, 219  
 Rational equation, 258  
   application, 265–67  
   in more than one variable, 261  
 Rational expression, 202  
 Rational expressions  
   applications of, 265–67  
   completely reduced, 208–10  
   definition of, 202  
   denominator of a, 203  
   difference of, 239, 246  
   domain of, 204  
   fundamental principle of, 207  
   least common denominator of, 242  
   numerator of, 203  
   product of, 233  
   quotient of, 235  
   sum of, 239, 246  
 Rationalizing a denominator, 379–81, 388, 427  
 Rational number, 27  
 Real number, 28  
 Real number line, 29  
 Reciprocal, 5, 94  
 Rectangular coordinate plane, 280  
 Reducing to lowest terms, 4, 208

Relation, 323  
   domain of a, 323  
   range of a, 323  
 Remainder, 5  
 Repeating decimal, 19  
 Root, 86  
 Root, extraneous, 260

**S**

Scientific notation, 148  
   computation using, 149  
   standard form of, 149  
 Set, 25  
 Set symbolism, 25  
 Signed numbers, 34  
 Similar terms, 80  
 Simple interest, 112  
 Simplifying rational expressions, 207  
 Slope, 297–98  
   definition of, 298–99  
   of a horizontal line, 300  
   of parallel lines, 309  
   of perpendicular lines, 310  
   undefined, 300–1  
   of a vertical line, 301  
 Slope-intercept form, 307  
 Solution, 86, 276  
   extraneous, 260  
 Solution set, 86, 187  
   of a linear equation in two variables, 277  
 Solutions of quadratic equations  
   by completing the square, 411–13  
   complex solutions of, 427  
   by extracting the roots, 405–6  
   by factoring, 404–5  
   by quadratic formula, 418–19  
 Special products, 135–37  
 Square of a binomial, 135  
 Square root, 367  
 Square root property, 405  
 Standard form of a quadratic equation, 404  
 Standard form of the equation of a line, 305  
 Statement, mathematical, 86  
 Straight line, 289  
 Strict inequality, 30  
 Subscripts, 75  
 Subset, 26  
 Substitution property, 72  
   method of solving a system of linear equations, 346–48  
 Subtraction, definition of, 41  
   decimals, 16  
 Subtraction, of rational expressions, 239  
 Subtraction of  
   fractions, 7–9, 239  
   more than two real numbers, 42  
   two real numbers, 41  
 Symbols  
   absolute value, 31  
   braces, 25, 42  
   brackets, 42  
   is an element of, 26  
   is approximately equal to, 19  
   is a subset of, 26  
   is greater than, 30  
   is greater than or equal to, 30

is less than, 30  
 is less than or equal to, 30  
 multiplication dot, 46  
 null set, 260  
 parentheses, 42  
 Symmetric property of equality, 89  
 Symmetry, 31  
 Systems of linear equations, 335  
   applications of, 351–55  
   consistent and independent, 337–38  
   dependent, 337–38  
   graphing of, 336–37  
   inconsistent, 337–38  
   linear, 335  
   solution by elimination, 340–44  
   solution by substitution, 346–48  
 Systems of linear inequalities, 360–61

**T**

Term, 67  
 Trinomial, 68  
 Trinomial, perfect square, 135, 409

**U**

Undefined, 53  
 Undefined slope, 301  
 Undirected distance, 31

**V**

Variable, 29, 67  
 Vertex of a parabola, 436–37  
 Vertical line, 293  
   slope of, 301

**W**

Weak inequality, 30  
 Whole numbers, 26

**X**

$x$ -axis, 280  
 $x$ -intercept, 290, 435

**Y**

$y$ -axis, 280  
 $y$ -intercept, 290, 435

**Z**

Zero, division involving, 53  
 Zero as an exponent, 142  
 Zero factor property, 47  
 Zero product property, 187





Free tunes.  
For Students. By Students.

Available for Fall 07 StudyBreaks  
[www.freeloadpress.com](http://www.freeloadpress.com)

Student musicians: Share your music. Make some change.

Learn more: [www.freeloadpress.com/studybreakmusic](http://www.freeloadpress.com/studybreakmusic)